

Application of Radioactive Fallout Cesium-137 for Measuring Soil Erosion and Sediment Accumulation Rates and Patterns: A Review

Jerry C. Ritchie* and J. Roger McHenry

ABSTRACT

Radioactive fallout ^{137}Cs (cesium-137) deposited across the landscape from atmospheric nuclear tests is strongly absorbed on soil particles limiting its movement by chemical and biological processes. Most ^{137}Cs movement in the environment is by physical processes; therefore, ^{137}Cs is a unique tracer for studying erosion and sedimentation. Cesium-137 loss from a watershed has been shown to correlate strongly with soil loss calculated by the Universal Soil Loss Equation (USLE) or measured from small runoff plates. By measuring spatial patterns of ^{137}Cs in vertical and horizontal planes across the landscape, rates of soil loss or deposition can be measured for different parts of a watershed. Even within landscape units, redistribution of soil can be mapped and erosion or deposition rates for different parts of individual fields measured and mapped. Sediment accumulation rates can be measured by comparing the vertical distribution of ^{137}Cs in sediments with the temporal deposition of fallout ^{137}Cs from the atmosphere to locate sediment horizons. Using these dated sediment horizons, sediment accumulation rates can be measured. Interpretations about the location of these dated horizons must consider particle size of the sediments, reworking of deposited sediments, diffusional movement of ^{137}Cs , and time rates of physical process in the sedimentation process. The ^{137}Cs technique can be used to determine sediment accumulation rates in a wide variety of depositional environments including reservoirs, lakes, wetlands, coastal areas, and floodplains. The bibliography shows that ^{137}Cs has been used widely for studying erosion and sedimentation in many different environments around the world.

SOIL EROSION and deposition of eroded soil particles in fields, floodplains, and water bodies are major environmental concerns around the world. Although soil erosion is a natural process, many human activities have increased rates of soil erosion. Soil erosion reduces soil productivity, scars the landscape, and causes downstream damage. Sediment accumulation in water bodies affects water quality, causes loss of storage capacity, affects biological activity, and reduces recreational potential. Better information is needed on sources, locations, and rates of erosion and on locations, patterns, and rates of deposition of eroded materials. Many qualitative and quantitative techniques have been developed and used to measure patterns and rates of erosion and sediment accumulation. Estimates of these patterns and rates have also been made using empirically and theoretically derived erosion models. However, few methods are capable of providing data on both erosion and sedimentation. Information on the relationship between erosion rates, delivery rates of eroded products, and sediment accumulation rates is needed if erosion and sedimentation is to be understood and monitored on a watershed.

J.C. Ritchie, USDA-ARS, Hydrology Lab., Beltsville, MD 20705; and J.R. McHenry, 7258 Spring Drive, Boulder, CO 80303. Received 27 Feb. 1989. *Corresponding author.

Published in J. Environ. Qual. 19:215-233 (1990).

Radioactive fallout ^{137}Cs has been used as a tracer to provide independent information on erosion and sedimentation rates and patterns. Sedimentation and sediment accumulation in this paper is used to indicate a vertical deposition of sediments (cm yr^{-1}). The purpose of this article is to review the development and application of a technique using radioactive fallout ^{137}Cs to measure rates and patterns of erosion and sedimentation. A bibliography is also provided.

WHY CESIUM-137?

Properties of radioactive ^{137}Cs make it unique as a tracer for studying erosion and sedimentation. There are no natural sources of ^{137}Cs . Cesium-137 is produced during nuclear fission. Thus, its presence in the environment is due to nuclear testing or to releases from nuclear reactors (Wise, 1980; Walling et al., 1986b). Cesium-137 was first released around sites used for early nuclear tests in 1945 (Carter and Moghissi, 1977). Widespread global dispersal of ^{137}Cs to the environment began with high-yield thermonuclear tests in November 1952 (Perkins and Thomas, 1980). In these tests, ^{137}Cs (and other radionuclides) was injected into the stratosphere where it circulates globally (Longmore, 1982). Cesium-137 moves from the stratosphere back to the troposphere and re-entry from the troposphere to the earth's surface (fallout) is strongly related to local precipitation patterns and rates (Davis, 1963; Longmore, 1982).

Temporal and spatial distribution of radioactive fallout on the earth's surface has been extensively measured and documented (Cambray et al., 1983; Larsen, 1985). Based on radioactive fallout measurements by the Health and Safety Laboratory (1972), initial input of ^{137}Cs into the global environment was estimated to be 1952 ± 2 yr (Robbins et al., 1978) with measurable amounts in soils generally beginning in 1954 (Wise, 1980; Longmore, 1982). Major periods of global deposition of ^{137}Cs fallout in 1958 and 1963/1964 and minor periods in 1971 and 1974 can be related to activity in aboveground nuclear testing (Carter and Moghissi, 1977). Periods of lower fallout rates can be related to moratoriums on testing (1958-1961) and the Test Ban Treaty of 1963. Because of the 1963 Test Ban Treaty, global radioactive fallout rates have decreased steadily except for minor periods in 1971 and 1974 caused by aboveground nuclear testing by non-treaty countries. The 1971 period was more significant in the southern hemisphere than in the northern hemisphere. Cambray et al. (1985) reported fallout deposition rates of ^{137}Cs in the northern hemisphere to be below their limits of detection in 1983 and 1984. Total fallout, as illustrated by ^{90}Sr (Fig. 1), is greater in the northern hemisphere than in the southern hemisphere (Larsen, 1985), because more atmospheric nuclear testing took place in the northern hemisphere. Local events, such as the Chernobyl accident (Volchok and Chieco, 1986), have significant impacts on regional




Fig. 1. Annual atmospheric fallout (^{90}Sr) in the northern and southern hemispheres. Cesium-137 is equal to 1.45 times ^{90}Sr at formation.

patterns of fallout radioactivity but have limited impacts on global fallout rates and patterns.

Cesium-137 is strongly adsorbed on clay and organic particles (Tamura and Jacobs, 1960; Schulz et al., 1960; Davis, 1963; Tamura, 1964a,b; Schulz, 1965; Brisbin et al., 1974) and is essentially nonexchangeable (Davis, 1963; Lomenick and Tamura, 1965). Adsorption on soils and sediments is rapid (Eyman and Kevern, 1975) with distribution in undisturbed soil profiles showing an exponential decrease with soil depth (Beck, 1966; Ritchie et al., 1970a, 1972b). Cesium-137 in plowed soils shows a uniform distribution through the plow layer (Cline and Rickard, 1972; Ritchie and McHenry, 1973a). Increased concentrations of competing ions such as Na, K (Coleman et al., 1963; Aston and Duursma, 1973), and H (Agre and Korogodin, 1960; Lerman and Lietzke, 1975) slightly decrease ^{137}Cs adsorption. However, Lomenick and Tamura (1965) removed less than 1% of adsorbed ^{137}Cs on soil and sediment samples using different acids and bases illustrating the strong adsorption between ^{137}Cs and soil particles and the limited mobility of ^{137}Cs by chemical processes.

Cesium-137 emits a strong gamma-ray (662 keV) making its measurement in environmental samples

relatively easy and accurate without special chemical preparation or separation (Ritchie and McHenry, 1973b; McCallan et al., 1980). Cesium-137 can be measured using a lithium-drifted germanium semiconductor detector or a thallium-activated sodium iodide crystal and a multichannel analyzer to separate and quantify the 662 keV gamma-ray peak from the gamma-ray spectrum. This specialized equipment is available in radiochemistry and physics laboratories.

With its long half-life (30 yr) and the total deposited fallout especially in the northern hemisphere (Cambray et al., 1983; Larsen, 1985), ^{137}Cs will be detectable in environmental samples for many years.

CESIUM-137 CYCLE ON THE LANDSCAPE

Although dry deposition is important locally around nuclear test sites, on a global scale, ^{137}Cs deposition from the atmosphere to the landscape (Fig. 2) has been found to be strongly related to rainfall with total fallout varying linearly with rainfall within latitudinal zones (Davis, 1963). Transport of ^{137}Cs across the landscape involves transfers between three primary components: vegetation, soil, and water.

Cesium-137 deposited on vegetation may be adsorbed or absorbed. Most adsorbed ^{137}Cs is washed from vegetation and moved to the soil (Davis, 1963; Dahlman et al., 1975). Rogowski and Tamura (1970a) found that 93% of the ^{137}Cs applied to grass washed off during the first year. Absorbed ^{137}Cs is released to soils when vegetation dies and decays. This turnover rate for absorbed ^{137}Cs depends on vegetation type and climate conditions. Uptake of ^{137}Cs by vegetation from soils (Fredrickson et al., 1958; Davis, 1963; Dahlman et al., 1975) or water (Eyman and Kevern, 1975; Garten et al., 1975) is low. Uptake is far less important than absorption from direct deposition on foliage in determining ^{137}Cs concentration in vegetation (Dahlman et al., 1975). Removal of ^{137}Cs from the landscape with harvested crops is very small (Brown et al., 1981a).

Cesium-137 in soils is due to direct deposition from the atmosphere, wash-off from vegetation, turnover from vegetation, redeposition of eroded soil particles, and deposition from water on floodplains and coastal regions. Loss by biological uptake from soils by vegetation can be considered negligible (Dahlman et al., 1975). Because ^{137}Cs is strongly adsorbed to cation exchange sites (Davis, 1963; Eyman and Kevern, 1975), movement in soils due to chemical or biological processes is limited (Tamura, 1964b; Wise, 1980; Campbell, 1982). Thus, physical processes, such as erosion and tillage, are major causes of redistribution of ^{137}Cs in soils and movement of ^{137}Cs from soils to water.

Cesium-137 in water is from direct deposition on water surfaces and input of ^{137}Cs adsorbed on eroded soil particles. Some fresh fallout ^{137}Cs may move from the landscape during large rainfall events before it becomes adsorbed by soils, but such movement of ^{137}Cs is very small (Brown et al., 1981a). Cesium-137 deposited on water surfaces will be adsorbed by suspended materials in water (Armstrong and Gloyne, 1969; Eyman and Kevern, 1975; Francis and Brinkley,




Fig. 2. Diagram of the ^{137}Cs cycle on the landscape.

1976). This suspended material is mostly eroded particles and becomes part of the sedimentation process. Except for overbank floods, which can redeposit ^{137}Cs on floodplains, deposited sediments in water bodies become the eventual sink for ^{137}Cs .

Although biological and chemical processes can move limited amounts of ^{137}Cs between vegetation, soil, and water, physical factors are the dominant processes transferring ^{137}Cs in natural systems (Davis, 1963; Rogowski and Tamura, 1970a,b; Ritchie et al., 1974; Dahlman et al., 1975). Because ^{137}Cs is uniformly distributed across the landscape and strongly adsorbed on soil particles, radioactive fallout ^{137}Cs can be used as a tracer for studying the physical processes of erosion and sedimentation. Thus, spatial measurements of ^{137}Cs can provide quantitative information on rates and patterns of erosion and sedimentation.

THE USE OF CESIUM-137 TO STUDY EROSION

Studies began in the early 1960s to measure the relationship between soil erosion and movement of fallout radionuclides. Menzel (1960) showed that loss of fallout ^{90}Sr from plots used to study soil erosion in Georgia and Wisconsin was related to measured soil loss. Graham (1963) found similar patterns for the movement of fallout ^{85}S and ^{131}I from soil erosion plots in Missouri. Frere and Roberts (1963) measured fallout ^{90}Sr loss from small research watersheds at Coshocton, OH, and found no loss of ^{90}Sr from grass-covered watersheds, whereas concentrations of ^{90}Sr in cultivated watersheds was only one-third to two-thirds that of the grass watersheds. They related this ^{90}Sr loss to measured soil loss from these watersheds. Dahlman and Auerbach (1968) found similar patterns for loss of ^{137}Cs applied to grass in Tennessee. Rogowski and Tamura (1965, 1970a,b) applied ^{137}Cs to small grassed test plots in Tennessee and found a logarithmic relationship between measured soil loss and ^{137}Cs loss. In 1974, a method was described for using measurements of the percentage loss of fallout ^{137}Cs to measure soil loss (Ritchie et al., 1974). In this article, fallout ^{137}Cs loss from different land uses in a watershed was shown to have a strong logarithmic relationship with soil loss estimated using the Universal Soil Loss Equation (USLE) (Wischmeier and Smith, 1978). Then they combined their results with those of Menzel (1960), Graham (1963), Frere and Roberts (1963), and Ro-




Fig. 3. Relationship between percent radionuclide loss and soil loss.

gowski and Tamura (1970a,b) to show a strong logarithmic relationship ($r = 0.94$) between soil loss in metric tons per hectare per year and percent radionuclide loss (Fig. 3). They concluded that soil erosion rates can be reliably measured by determining the percent of fallout ^{137}Cs loss from soils.

McHenry and Ritchie (Ritchie et al., 1971, 1974, 1982; McHenry and Ritchie, 1977a; McHenry et al., 1978; Mitchell et al., 1980; McHenry and Bubenzer, 1985; Spomer et al., 1985) began a series of studies in the early 1970s using the spatial distribution of ^{137}Cs in watersheds to determine (i) patterns of soil erosion and redeposition of eroded material within the watershed, (ii) movement of eroded material from watersheds, and (iii) deposition of eroded material in water bodies. These studies showed that ^{137}Cs measured at noneroded sites was equal to the fallout ^{137}Cs deposited in the area and was in the upper part of the soil profile or in the plow layer of cultivated soils and thus

could be used as the baseline for the input of ^{137}Cs to an area. At sites where soil erosion was occurring, total ^{137}Cs in the soil profile was less than the input from fallout. At deposition sites in the watershed, total ^{137}Cs was greater than input from fallout and/or the depth distribution of ^{137}Cs in soil profiles was greater than that measured in a noneroded site or deeper than the plow layer. McHenry and Ritchie (1977a) concluded that by measuring spatial patterns of ^{137}Cs in both vertical and horizontal planes across the landscape, rates of soil loss or deposition could be measured for different parts of a watershed. Even within landscape units, redistribution of soil could be mapped and erosion or deposition rates for different parts of individual fields measured and mapped from measurements of the spatial distribution of fallout ^{137}Cs . Erosion and deposition rates measured using the spatial distribution of ^{137}Cs were related to slope steepness, shape, and length. Flat areas at the top of slopes usually showed little soil loss. Flat areas at the base of slopes and concave slopes in fields often showed deposition. Thus measurements of the spatial distribution of ^{137}Cs could be used to show erosion patterns in fields, the redistribution of soils within fields, and soil loss from fields.

Martz and DeJong (1987) emphasized that the ^{137}Cs technique measures NET soil loss at each site rather than the total soil loss estimated by other methods and thus provides better information on true erosion rates and sediment delivery. Net soil loss was defined as soil loss minus soil deposition at a site. Other researchers have used ^{137}Cs in a wide variety of landscape and environmental conditions to measure soil loss, soil redeposition, and to develop budgets for erosion and sedimentation (i.e., Wise, 1977, 1980; McCallan et al., 1980; Brown et al., 1981a,b; Loughran et al., 1981, 1982, 1986a,b,c, 1987a,b; Campbell et al., 1982, 1985, 1986a,b; DeJong et al., 1982, 1983, 1986; Longmore et al., 1983a,b; Elliott et al., 1984; Kachanoski and DeJong, 1984; Kiss et al., 1986; Lance et al., 1986a,b,c; Walling et al., 1986a,b; Berghe and Gulinck, 1987; Kachanoski, 1987; Martz and DeJong, 1987; McIntyre et al., 1987; Menzel et al., 1987; Pennock and DeJong, 1987; Lowrance et al., 1988).

To use fallout ^{137}Cs to measure erosion, a baseline input of ^{137}Cs must be determined. This baseline input to the environment is due to fallout from the atmospheric nuclear testing rather than natural sources (Davis, 1963; Wise, 1980) and its initial distribution across the landscape is uniform or can be related to measurable environmental gradients (i.e., precipitation). It is not critical to have direct measurements of the actual ^{137}Cs deposited in a watershed, because local input of ^{137}Cs to a landscape unit can be estimated by measuring ^{137}Cs in soil profiles where neither erosion nor soil deposition has occurred (Ritchie et al., 1971, 1974; Campbell, 1982; Walling et al., 1986a,b; Martz and DeJong, 1987). It is important to assume a uniform deposition pattern across the landscape unit being studied or to be able to estimate the distribution pattern across a large watershed based on an environmental gradient so that measurements made at non-eroding sites can be used to determine fallout input to the study area.

Another technique for determining baseline meas-

urements is to develop a monitoring program and measure actual changes in ^{137}Cs with time at each site (Kachanoski and DeJong, 1984; Kachanoski, 1987). Such a monitoring program allows a comparison of actual measurements rather than having to make assumptions about uniform fallout patterns. Kachanoski (1987) has shown high variability in the spatial distribution of ^{137}Cs on landscape units.

Comparisons between baseline and current measurements of the vertical and horizontal distribution of ^{137}Cs across the landscape will reflect the movement of ^{137}Cs adsorbed on soil particles within and between soils and water (Fig. 2) by the physical process of erosion. Movement of ^{137}Cs in soils by biological and chemical processes is small (Tamura, 1964a,b) in comparison with movement of ^{137}Cs by physical processes (i.e., water and wind) that transport soil particles across the landscape. Loss of ^{137}Cs from a site is strongly related to the amount of soil loss (Ritchie et al., 1974; DeJong et al., 1982; Martz and DeJong, 1987). Thus, the spatial distribution of ^{137}Cs will reflect the pattern of atmospheric fallout as it has been modified by patterns of erosion and deposition of soil particles. McHenry and Bubenzer (1985) measured the spatial distribution of ^{137}Cs in a Wisconsin watershed and estimated that 50 to 100 times more soil was redistributed within a field than was removed from the field by erosion.

Walling et al. (1986b) incorporated data on input, redistribution, and output of ^{137}Cs from a watershed to measure erosion and sedimentation on a watershed with a single technique. Such watershed budgets of fallout ^{137}Cs (Ritchie et al., 1971, 1974, 1982; Ritchie and McHenry, 1976; McHenry and Ritchie, 1975; Bazzoffi and Panicucci, 1983; DeJong et al., 1983; Walling et al., 1986b) and studies of the spatial distribution of ^{137}Cs provide a unique source of data for studying and understanding erosion and sedimentation.

THE USE OF CESIUM-137 TO STUDY SEDIMENT ACCUMULATION

In 1961, Ravera (1961) published a report on the buildup of fallout radioactivity in sediments of Lake Maggiore in Italy. This report included a vertical profile of the radioactive beta-ray activity in the sediment. Later, using this sediment profile, Ravera and Premazzi (1971) reported that "the sedimentation rate was evaluated by knowing the distribution in time of the fall-out in air."

Studies in the mid 1960s by Pickering et al. (1965, 1966); Carrigan and Pickering (1967); Pickering (1969) at Oak Ridge, TN, on the Clinch River; and Nelson et al. (1966) at Hanford, WA, on the Columbia River, related the vertical distribution of total gamma activity, total beta activity, ^{137}Cs , and other radionuclides in sediments to the time history of radionuclide release from Oak Ridge or Hanford nuclear processing facilities. These studies were based on relatively high levels of radioactivity compared with levels of radioactivity in the environment due to radioactive fallout from nuclear tests.

In 1968, Schreiber (1968) and Schreiber et al. (1968) related the vertical distribution of total beta counts in marine sediments from the North Adriatic Sea to at-

atmospheric radioactive fallout rates from nuclear testing. They postulated that the vertical distribution of total beta activity in sediments was related to yearly fallout rates and thus could be used to date horizons in sediment profiles. They attributed most of the radioactivity measured to fallout ^{144}Ce and ^{90}Sr . Independent measurements of sediment accumulation rates for these marine sediment profiles by other methods were not available for comparison with the rates measured from the beta radioactivity profiles.

In the early 1970s, published reports of independent studies in the USA (Ritchie et al., 1970c,d, 1973; Robbins and Edgington, 1972, 1975; McHenry et al., 1973), Italy (Ravera and Premazzi, 1971), Switzerland (Krishnaswami et al., 1971), England (Pennington et al., 1973), and Israel (Stiller and Assaf, 1973) concluded that the vertical distribution of fallout ^{137}Cs in sediments could be related to the time distribution of radioactive fallout ^{137}Cs deposition from the atmosphere and thus used to infer a geochronology for sediment profiles. These studies measured sediment accumulation rates and patterns by measuring the vertical distribution of ^{137}Cs in sediment profiles. Ritchie et al. (1973) and Ritchie and McHenry (1975a) compared their sediment accumulation rates (cm yr^{-1}) measured from ^{137}Cs in sediment profiles with sediment accumulation rates measured at the same site by standard sediment survey methods (Fig. 4) and showed that comparable sedimentation rates were measured by the different techniques.

To use ^{137}Cs to measure sediment accumulation rates and patterns, the temporal pattern of fallout from atmospheric nuclear testing or release from nuclear facilities must be known (Davis, 1963; Wise, 1980). A comparison of these temporal measurements of the input of atmospheric fallout ^{137}Cs to the environment with measurements of ^{137}Cs in sediment profiles can

be used to date different horizons in sediment profiles. By knowing the depth of these different ^{137}Cs horizons, rates of sediment accumulation can be calculated.

Cesium-137 input to sediment profiles is from direct deposit at the site or from the movement of ^{137}Cs deposited on the upstream watershed by the physical processes that transport eroded particles. In water bodies, ^{137}Cs deposited on the water surface is adsorbed to suspended materials in the water (Armstrong and Gloyna, 1969; Eyman and Kevern, 1975; Francis and Brinkley, 1976). This suspended material, which is primarily eroded soil particles also carrying ^{137}Cs , will be deposited as part of the sediment deposition process. Most of the suspended material entering a water body is deposited rapidly; thus, the sequential buildup of deposited sediment will reflect the time distribution pattern of fallout ^{137}Cs .

Because ^{137}Cs is strongly adsorbed on soil particles in the surface soil layer (Beck, 1966) and eroded materials tend to come from the same area within a watershed each year (Sprayberry and Bowie, 1969), ^{137}Cs concentration on soil particles being eroded from a watershed and moved to depositional sites will reflect the temporal fallout of ^{137}Cs .

Dates for two sediment horizons (1954 and 1964) can be identified in most deposition sites in the northern hemisphere based on the measurement of ^{137}Cs concentration. These horizons may not be as definitive in the southern hemisphere due to lower rates of fallout (Longmore, 1982). Sediment horizons for 1958 can sometimes be identified. A 1971 sediment horizon has been identified in the southern hemisphere (Campbell, 1983) and tentatively identified in a few northern hemisphere sediment profiles (McHenry et al., 1984). The Chernobyl accident has labeled a sediment horizon in much of Europe (Buesseler et al., 1987). Thus, with several depositional horizons marked by ^{137}Cs , com-




Fig. 4. Depth distribution of ^{137}Cs in sediment profiles from Mule Creek B reservoir in Iowa. Dates on the right of each profile indicate

parison of sediment accumulation rates can be made for different time periods from measurements of a single sediment profile.

The ^{137}Cs technique has some advantages over conventional techniques of measuring sediment accumulation rates. Measurements of sediment accumulation rates can be made rapidly from measurements of a single sediment profile with no requirement to return to the site to make repeated measurements. Interpretations about sediment accumulation rates are made by comparing the vertical distribution of ^{137}Cs in sediments with the known time deposition of fallout ^{137}Cs from the atmosphere. This time distribution (monthly and annual) of radioactive fallout has been extensively measured and documented for different areas around the world (Cambray et al., 1983; Larsen, 1985).

There are certain factors about the ^{137}Cs technique that must be considered when interpretations of the vertical distribution of ^{137}Cs in sediments are used to measure sediment accumulation rates. Only two dates (1954 and 1964) can be determined for most sediment profiles. If more detail on the time rate of sediment accumulation is needed, other techniques should be used.

Determining ^{137}Cs near the detection limits can cause inaccuracies in locating the 1954 sediment horizon. Radioactive decay of ^{137}Cs (half-life = 30 yr) continues to reduce the amount of ^{137}Cs in the environment and thus limits the detection of the 1954 horizon (Campbell, 1982) due to the small amount of ^{137}Cs originally present in the 1954 horizon. By 1984, only half of the original amount of ^{137}Cs deposited in the 1954 sediment horizon would be left. This is especially critical in the southern hemisphere where fallout rates are lower than in the northern hemisphere (Fig. 1).

Bulk density of the sediment sample must be known to be able to calculate total ^{137}Cs . Also, because ^{137}Cs is strongly adsorbed on clay-size particles, knowledge about the particle size distribution within the sediment profile is helpful in interpretation (Fig. 5). An increase

in the amount of sand-size particles in sediment profiles will cause an apparent decrease in the concentration of ^{137}Cs at that point in the sediment profile that cannot be related to any time change in atmospheric fallout rates of ^{137}Cs . For some sediment profiles it is probably better to express ^{137}Cs concentration in units per unit of clay to account for the differences in ^{137}Cs distribution due to differences in particle size distribution in the sediment profile. In depositional environments (i.e., barrier islands) where sands dominate the sediment profile, ^{137}Cs will be of limited value in determining sedimentation.

A time lag exists between the time of atmospheric deposition of ^{137}Cs and the time of deposition of ^{137}Cs in sediment profiles. This time lag is from factors controlling the processes of sediment deposition. Early research in a small watershed indicated a time lag of 6 to 12 mo between the time of atmospheric fallout and the time of ^{137}Cs deposition in a reservoir sediment (Ritchie et al., 1973). However, the actual time lag will depend on the rates and frequency of different natural processes in each water body and watershed (Wise, 1980). Cesium-137 may even help understand these time lag processes (Walling et al., 1986a). This time lag should be considered very carefully in assigning dates to sediment horizons.

Physical reworking of sediments by animals, wave action, or other methods redistribute ^{137}Cs within the profile and thus spread ^{137}Cs over a larger part of the profile and reduce maximum concentrations of ^{137}Cs . A number of studies have reported this redistribution (i.e., Robbins and Edgington, 1975; Robbins et al., 1977b, 1979; Hakanson and Kallstrom, 1978; Krezoski et al., 1978; Krezoski and Robbins, 1985; Fisher et al., 1980; Kacieszczenko and Banasik, 1981; Demaster et al., 1985; Robbins, 1986a; Anderson et al., 1987; Sharma et al., 1987). Kacieszczenko and Banasik (1981) used a model to predict the effects of bioturbation on ^{137}Cs movement and showed that movement can occur. In general, any reworking of sediment broadens the ^{137}Cs peaks and makes interpretation more difficult. However, such movement will not likely change the position of the major ^{137}Cs horizons (Pennington et al., 1973, 1976; Ritchie et al., 1973; Krishnaswami et al., 1971; Robbins and Edgington, 1975; Wise, 1980; Campbell, 1982). Such movement, however, can have a major effect on the position of the ^{137}Cs horizon deposited in 1954. By spreading detectable levels of ^{137}Cs into lower parts of the profile, the first detectable level of ^{137}Cs would be lower than it should be. Localized reworking of the near-surface sediments can also displace ^{137}Cs horizons, especially in areas with low sediment deposition rates. In general, the lower the sediment accumulation rate the greater effect reworking of sediments will have on the redistribution of ^{137}Cs in sediment profiles.

Diffusional movement of ^{137}Cs also redistribute ^{137}Cs in sediment profiles. Although diffusional movement is usually limited (Davis, 1963; Tamura, 1964a,b; Aston and Duursma, 1973; Lomenick and Tamura, 1965), it does occur and can be a problem in certain environments (i.e., Krishnaswami et al., 1971; Evans et al., 1983; Davis et al., 1984; Torgenson and Longmore, 1984). Again, such movement will not likely change the position of ^{137}Cs horizons (Penning-




Fig. 5. Depth distribution of ^{137}Cs in a sediment profile from Tortugas Arroyo no. 1 reservoir in New Mexico compared with particle size distribution in the same profile.

ton et al., 1973, 1976; Ritchie et al., 1973; Krishnaswami et al., 1971; Robbins and Edgington, 1975; Wise, 1980; Campbell, 1982). However, for soft-water lakes (Davis et al., 1984), diffusion of ^{137}Cs can be a major problem. Diffusional movement of ^{137}Cs has the greatest effect on the location of the ^{137}Cs horizon deposited in 1954; thus, care needs to be taken in determining the location of this horizon.

Cesium-137 is difficult to use in water bodies with low sediment accumulation rates ($< 1 \text{ cm yr}^{-1}$) because of sampling problems. If sediment accumulation is low, then sampling of horizons in the detail necessary to determine ^{137}Cs horizons in the profile may be difficult.

The ^{137}Cs technique can be used only with sediments deposited since 1954, the year that radioactive fallout ^{137}Cs was first deposited in measurable amounts. Also, as noted earlier, determining the position of the 1954 horizon will become more difficult with time due to the natural decay of ^{137}Cs . Thus, one is limited to determining sediment accumulation rates since 1954. Other techniques (e.g., ^{210}Pb , ^{14}C , pollen) can be used to measure older deposits.

Fallout ^{137}Cs has been declining in the environment since 1964; thus, the last major sediment ^{137}Cs marker is usually 1964. For determinations of sediment accumulation rates after 1964 or sediment accumulation rates in a reservoir built since 1964, the ^{137}Cs technique may not be applicable. There are exceptions to this, such as a horizon labeled by fallout from the Chernobyl accident in Europe or other similar events.

SUMMARY

Research beginning in the late 1960s has shown that ^{137}Cs deposited across the landscape by fallout from nuclear test is a unique and useful tool for studying erosion and sedimentation. Cesium-137 has been used to measure erosion rate and delineate patterns of erosion and deposition within landscape units to provide data on net erosion from a site. These data provide a basis for determining budgets and patterns of soil redistribution within a field and actual soil loss from the field. Cesium-137 has also been used to determine the sediment accumulation rates in a wide variety of depositional environments for sediment deposited since 1954. By providing data on erosion rates and patterns and on sediment accumulation rates and patterns, ^{137}Cs is a unique tool for studying the complete erosion and sedimentation cycle across the landscape.

ACKNOWLEDGMENTS

Jeanne Morrisette, Arlene Keller, Leoma Kruzic, and Mary Jane Evans assisted in the preparation of the bibliography.

The References section contains publications and reprints collected by the authors over the time that they have been involved in research on the use of fallout ^{137}Cs to study erosion and sedimentation. In our first paper on ^{137}Cs (Ritchie et al., 1970b), we stated that our purpose was "to describe the use of radioactive fallout, as ^{137}Cs , as a tracer of erosion, sediment transport, and sediment deposition within a watershed." Our research and that of our colleagues have more than met this purpose as stated at the meeting of the Mississippi Water Resources Conference on 15 Apr. 1970.

Our research began in 1968 as a cooperative research effort between the USDA-ARS, Sedimentation Laboratory in Ox-

ford, MS, and the United States Atomic Energy Commission (now the U.S. Department of Energy). We would like to thank all colleagues, associates, co-workers, and friends who over the years have provided us support, labor, ideas, criticisms, and encouragement.

REFERENCES¹

- Ackermann, F., H. Bergmann, and U. Schleichert. 1983. Monitoring of heavy metals in coastal and estuarine sediments—A question of grain-size: 20 μm versus 60 μm . *Environ. Technol. Lett.* 4:317–328.
- †Agre, A.L., and V.I. Korogodin. 1960. The distribution of radioactive contamination in a stagnant reservoir. *Med. Radiol.* 5:161–175.
- Alberts, J.J., and R.N. Muller. 1979. The distribution of $^{239,240}\text{Pu}$, ^{238}Pu , and ^{137}Cs in various particle size classes of Lake Michigan sediments. *J. Environ. Qual.* 8:20–22.
- Alberts, J.J., L.J. Tilly, and T.J. Vigerstad. 1979. Seasonal cycling of cesium-137 in a reservoir. *Science (Washington, DC)* 203:649–651.
- Alberts, J.J., and M.A. Wahlgren. 1981. Concentrations of $^{239,240}\text{Pu}$, ^{137}Cs , and ^{90}Sr in the waters of the Laurentian Great Lakes. Comparison of 1973 and 1976 values. *Environ. Sci. Technol.* 15:94–98.
- Aleksan'yan, O.M. 1978. A study of the vertical distribution of ^{90}Sr and ^{137}Cs in bottom sediments of a sea in Azov USSR. (In Russian.) *Ekologiya* 5:95–97.
- Aleksan'yan, O.M., and Y.Y. Katsnel'son. 1979. Radioactive pollution of bottom sediments and its prevention. *Sov. J. Ecol.* 10:243–245.
- Alhonen, P. 1979. The sedimentary record of the cultural eutrophication and pollution of lakes in Finland. *Arch. Hydrobiol.* 86:13–26.
- Aliyev, D.A., M.A. Abdullayev, and A.T. Tadiyev. 1977. Distribution of Sr^{90} and Cs^{137} in the plowed soils of the dry (subtropical) steppe zone of the lesser Caucasus in the Azerbaijan SSR. *Pochvovedeniye* 7:34–35.
- Aldredge, A.W., and F.W. Whicker. 1972. A method for measuring soil erosion and deposition with beta particle attenuation. *J. Range Manage.* 25:393–398.
- †Anderson, R.F., S.L. Schiff, and R.H. Hesslein. 1987. Determining sediment accumulation and mixing rates using ^{210}Pb , ^{137}Cs , and other tracers: Problems due to postdepositional mobility or coring artifacts. *Can. J. Fish. Aquat. Sci.* 44:231–250.
- Anspaugh, L.R., R. J. Catlin, and M. Goldman. 1988. The global impact of the Chernobyl reactor accident. *Science (Washington, DC)* 242:1513–1519.
- Appleby, P.G., J.A. Dearing, and F. Oldfield. 1985. Magnetic studies of erosion in a Scottish lake catchment. 1. Core chronology and correlation. *Limnol. Oceanogr.* 36:1144–1153.
- Appleby, P.G., and F. Oldfield. 1979. Correspondence on Edgington and Robbins (1976). *Environ. Sci. Technol.* 13:478–480.
- Appleby, P.G., and F. Oldfield. 1983. The assessment of ^{210}Pb data from sites with varying sediment accumulation rates. *Hydrobiologia* 103:29–35.
- Appleby, P.G., F. Oldfield, R. Thompson, and P. Huttunen. 1979. ^{210}Pb dating of annually laminated lake sediments from Finland. *Nature (London)* 280:53–55.
- Armentano, T.V. 1975. Sedimentation rates in a Long Island marsh determined by ^{210}Pb dating. *Limnol. Oceanogr.* 20:452–456.
- †Armstrong, N.E., and E.F. Gloyna. 1969. Mathematical models for the dispersion of radionuclides in aquatic systems. p. 329–335. In D.J. Nelson and F.C. Evans (ed.) *Symposium on radioecology*, Ann Arbor, MI. 15–17 May 1967. USAEC, Doc. CONF-670503. USAEC, Washington, DC.
- Ashley, G.M. 1977. Sedimentology of a tidal river, Pitt River, B.C. Ph.D. thesis. Univ. of British Columbia, Vancouver, BC.
- Ashley, G.M., and L.E. Moritz. 1979. Determination of lacustrine sedimentation rates by radioactive fallout (^{137}Cs), Pitt Lake, British Columbia. *Can. J. Earth Sci.* 16:965–970.
- Ashwood, T.L., and C.R. Olsen. 1988. Pearl Harbor bombing attack: A contamination legacy revealed in the sedimentary record. *Mar. Pollut. Bull.* 19:68–71.
- †Aston, S.R., and E.K. Duursma. 1973. Concentration effects on ^{137}Cs , ^{65}Zn , ^{60}Co , and ^{106}Ru sorption by marine sediments, with geochemical implication. *Neth. J. Sea Res.* 6:225–240.
- Aston, S.R., and J.E. Rae. 1982. The deposition of windscale radiocaesium in the Wyre estuary and the measurement of sedimentation rates. *Mar. Environ. Res.* 7:83–90.
- Aston, S.R., and D.A. Stanners. 1979. The determination of es-

¹ References cited in the text are designated by †. The other citations are relevant to this subject area.

- tuarine sedimentation rates by $^{134}\text{Cs}/^{137}\text{Cs}$ and other artificial radionuclide profiles. *Estuarine Coastal Mar. Sci.* 9:529-541.
- Aston, S.R., and D.A. Stanners. 1982. Local variability in the distribution of windscale fission products in estuarine sediments. *Estuarine Coastal Shelf Sci.* 14:167-174.
- Bachhuber, H., K. Bunzl, and W. Schimmack. 1986. Spatial variability of distribution of ^{137}Cs , ^{65}Zn , ^{57}Co , ^{109}Cd , ^{141}Ce , ^{103}Ru , ^{95}mTc , and ^{131}I in a cultivated soil. *Nucl. Technol.* 72:359-371.
- Bachhuber, H., K. Bunzl, and W. Schimmack. 1987. Spatial variability of fallout ^{137}Cs in the soil of a cultivated field. *Environ. Monit. Assess.* 8:93-101.
- Bachhuber, H., K. Bunzl, W. Schimmack, and I. Gans. 1982. The migration of ^{137}Cs and ^{90}Sr in multilayer soils: Results from batch, column, and fallout investigations. *Nucl. Technol.* 59:291-301.
- Barbeau, C., R. Bougie, and J.E. Cote. 1981. Variations spatiales et temporelles du césium-137 et du carbone dans des sédiments du fjord du Saguenay. (In French.) *Can. J. Earth Sci.* 18:1004-1011.
- Battarbee, R.W. 1978. Observations on the recent history of Lough Neagh and its drainage basin. *Philos. Trans. R. Soc. London* 281:303-345.
- Battarbee, R.W. 1986. The eutrophication of Lough Erne inferred from changes in diatom assemblages of ^{210}Pb and ^{137}Cs -dated sediment cores. *Proc. R. Ir. Acad. Sec. B*: 86:141-168.
- Battarbee, R.W., G. Digerfeldt, P. Appleby, and F. Oldfield. 1980. Palaeoecological studies of the recent development of Lake Vaxjö. III. Reassessment of recent chronology on the basis of modified ^{210}Pb dates. *Arch. Hydrobiol.* 89:440-446.
- Battiston, G.A., S. Degetto, R. Gerbas, and G. Sbrignadello. 1989. Determination of sediment composition and chronology as a tool for environmental impact investigations. *Mar. Chem.* 26:91-100.
- Battiston, G.A., S. Degetto, R. Gerbas, G. Sbrignadello, and L. Tositti. 1988. The use of ^{210}Pb and ^{137}Cs in the study of sediment pollution in a lagoon of Venice. *Sci. Total Environ.* 77:15-23.
- Baxter, M.S., R.W. Crawford, D.S. Swan, and J.G. Farmer. 1981a. ^{210}Pb dating of a Loch Lomond sediment core by conventional and particle track methods and some geochemical observations. *Earth Planet. Sci. Lett.* 53:434-444.
- Baxter, M.S., J.G. Farmer, I.G. McKinley, D.S. Swan, and W. Jack. 1981b. Evidence of the unsuitability of gravity coring for collecting sediment in pollution and sedimentation rate studies. *Environ. Sci. Technol.* 15:843-846.
- Baxter, M.S., I.G. McKinley, A.B. MacKenzie, and W. Jack. 1979. Windscale radiocaesium in the Clyde Sea area. *Mar. Pollut. Bull.* 10:116-120.
- †Bazzoffi, P., and M. Panicucci. 1983. Erosione sui versanti e conseguente sedimentazione in piccoli serbatoi artificiali. *Istituto. (In Italian.) Sperimentale Studio e Difesa Suolo, Firenze* 14:127-178.
- Beasley, T.M., R. Carpenter, and C.D. Jennings. 1982. Plutonium, ^{241}Am and ^{137}Cs ratios, inventories and vertical profiles in Washington and Oregon continental shelf sediments. *Geochim. Cosmochim. Acta* 46:1931-1946.
- Beasley, T.M., and C.D. Jennings. 1984. Inventories of 239 , ^{240}Pu , ^{241}Am , ^{137}Cs , and ^{60}Co in Columbia River sediments from Hanford to the Columbia River estuary. *Environ. Sci. Technol.* 18:207-212.
- †Beck, H.L. 1966. Environmental gamma radiation from deposited fission products 1960-1964. *Health Phys.* 12:313-322.
- Bell, G.L., and B.J. Eadie. 1983. Variations in the distribution of suspended particles during an upwelling event in Lake Michigan in 1980. *J. Great Lakes Res.* 9:559-567.
- Belperio, A.P., B.W. Smith, H.A. Polach, C.A. Nittrouer, D.J. DeMaster, J.R. Prescott, J.R. Hails, and V.A. Gostin. 1984. Chronological studies of the Quaternary marine sediments of northern Spencer Gulf, South Australia. *Mar. Geol.* 61:265-296.
- Benninger, L.K., and M.M. Brinson. 1988. Chronometric radionuclides in sediments of Cedar Island Salt Marsh, North Carolina. *EOS* 69:379.
- Berg, A., M. Merlini, O. Ravera, and V. Tonolli. 1961. Accumulation of fission products from fall-out in Lake Biota (Lago Maggiore). Final Rep. of the I.A.E.A. Contract 59. Int. Atomic Energy Agency, Vienna.
- Bergersen, E.P. 1987. Aldrin, dieldrin, and mercury profiles in recent lake sediments at the Rocky Mountain Arsenal, Colorado. *Arch. Environ. Contam. Toxicol.* 16:61-67.
- †Berghe, I.V., and H. Gulink. 1987. Fallout ^{137}Cs as a tracer for soil mobility in the landscape framework of Belgian loamy region. *Pedologie* 37:5-20.
- Bertine, K.K., and M.F. Mendeck. 1978. Industrialization of New Haven, Conn., as recorded in reservoir sediments. *Environ. Sci. Technol.* 12:201-207.
- Bickford, M.E., and W.R. Van Schmus. 1979. Geochronology and radiogenic isotope research. *Rev. Geophys. Space Phys.* 17:824-839.
- Bloemendal, J., F. Oldfield, and R. Thompson. 1979. Magnetic measurements used to assess sediment influx at Llyn Goddionduon. *Nature (London)* 280:50-53.
- Bodbacka, L. 1986. Sediment accumulation in Lake Lilla Ullfjärden and Stora Ullfjärden, Sweden. *Hydrobiologia* 143:337-342.
- Bonnett, P.J.P. 1989. A review of the erosional behavior of radionuclides in selected drainage basins. AERE-R-13631. U.K. AERE, Harwell, U.K.
- Bonny, A.P. 1976. Recruitment of pollen to the seston and sediment of some Lake District lakes. *J. Ecol.* 64:859-887.
- Bonny, A.P. 1978. The effect of pollen recruitment processes on pollen distribution over the sediment surface of a small lake in Cumbria. *J. Ecol.* 66:385-416.
- Bopp, R.F., H.J. Simpson, C.R. Olsen, and N. Kostyk. 1981. Polychlorinated biphenyls in sediments of the tidal Hudson River, New York. *Environ. Sci. Technol.* 15:210-216.
- Boudreau, B. 1986a. Mathematics of tracer mixing in sediments: I. Spatially-dependent, diffusive mixing. *Am. J. Sci.* 286:161-198.
- Boudreau, B. 1986b. Mathematics of tracer mixing in sediments: II. Nonlocal mixing and biological conveyor-belt phenomena. *Am. J. Sci.* 286:199-238.
- Bowen, V.T., V.E. Noshkin, and H.L. Volchok. 1970. Can land runoff be a major vector of fallout to the oceans? USAEC Rep. HASL-217, p. 1-119-125. USAEC Health and Safety Lab., New York.
- †Brisbin, I.L., Jr., R.J. Beyers, R.W. Dapson, R.A. Geiger, J.B. Gentry, J.W. Gibbons, M.H. Smith, and S.K. Woods. 1974. Patterns of radiocesium in the sediments of a stream channel contaminated by production reactor effluents. *Health Phys.* 27:19-27.
- Broom, M.J., P.D. Grimwood, and E.G. Bellinger. 1973. Caesium-137: Its accumulation in a littoral community. *Mar. Pollut. Bull.* 6:24-26.
- Brown, R.B. 1980. Agricultural erosion and sediment in the Western Willamette Valley as indicated by the redistribution of cesium-137. Ph.D. thesis. Oregon State Univ., Corvallis (DAI no. 41:1191-B).
- †Brown, R.B., N.H. Cutshall, and G.F. King. 1981a. Agricultural erosion indicated by ^{137}Cs redistribution: I. Level and distribution of ^{137}Cs activity in soils. *Soil Sci. Soc. Am. J.* 45:1184-1190.
- Brown, R.B., and J.S. Hickman. 1980. Runoff-contributing areas in fall-planted fields. p. 138-144. In M.E. Harward et al. (ed.) Reports on erosion, sediment, and water quality in the high winter rainfall zone of the northwestern United States. Oregon Agric. Exp. Stn. Spec. 602.
- Brown, R.B., and G.F. Kling. 1980. Erosion and sediment as indicated by redistribution of cesium-137. p. 90-108. In Oregon Agric. Exp. Stn. Spec. Rep. 602.
- †Brown, R.B., G.F. Kling, and N.H. Cutshall. 1981b. Agricultural erosion indicated by ^{137}Cs redistribution: II. Estimates of erosion rates. *Soil Sci. Soc. Am. J.* 45:1191-1197.
- Brugam, R.B. 1978. Pollen indicators of land-use change in southern Connecticut. *Quat. Res. (NY)* 9:349-362.
- Brunskill, G.J., S.D. Ludlam, and T.H. Peng. 1984. Fayetteville Green Lake, New York, U.S.A. VIII. Mass balance for ^{137}Cs in water, varved and non-varved sediments. *Chem. Geol.* 44:101-117.
- Brush, G.S. 1984. Patterns of recent sediment accumulation in Chesapeake Bay (Virginia-Maryland, U.S.A.) tributaries. *Chem. Geol.* 44:227-242.
- Brush, G.S., E.A. Martin, R.S. DeFries, and C.A. Rice. 1982. Comparisons of ^{210}Pb and pollen methods for determining rates of estuarine sediment accumulation. *Quat. Res. (NY)* 18:196-217.
- Bubbenzer, G.D. 1986. Quantifying erosion losses using cesium-137. Paper of the 1986 Summer Meetings of the American Society of Agricultural Engineers (ASAE Microfiche 86-2042).
- †Buesseler, K.O., H.D. Livingston, S. Honjo, B.J. Hay, S.J. Mangani, E. Degens, V. Ittekkot, E. Izdar, and T. Konuk. 1987. Chernobyl radionuclides in a Black Sea sediment trap. *Nature (London)* 329:825-828.
- Buesseler, K.O., and E.R. Sholkovitz. 1987. The geochemistry of fallout plutonium in the North Atlantic: I. A pore water study in shelf, slope and deep-sea sediments. *Geochim. Cosmochim. Acta* 51:2605-2622.
- Bunzl, K., W. Schimmack, K. Kreutzer, and R. Schierl. 1989. The migration of fallout ^{134}Cs , ^{137}Cs , and ^{106}Ru from Chernobyl and ^{137}Cs from weapons testing in a forest. *Z. Pflanzenenernähr. Bodenk.* 152:39-44.
- Bunzl, K., and W. Schultz. 1985. Distribution coefficients of ^{137}Cs and ^{85}Sr by mixtures of clay and humic material. *J. Radioanal. Nucl. Chem.* 90:23-37.
- Bush, B., L.A. Shane, M. Wahlen, and M.P. Brown. 1987. Sedimentation of 74 PCB congeners in the upper Hudson River. *Chemosphere* 16:733-744.
- Cahill, R.A., and J.D. Steele. 1986. ^{137}Cs as a tracer of recent sed-

- imentary processes in Lake Michigan. *Hydrobiologia* 143:29-35.
- Calcagno, T.H., and G.M. Ashley. 1984. Sediment processes in an impoundment, Union Lake, New Jersey. *Environ. Geol. Water. Sci.* 6:237-246.
- †Cambray, R.S., K. Playford, and N.J. Lewis. 1983. Radioactive fallout in air and rain: Results to the end of 1982. U.K. Atomic Energy Authority Rep. AERE-R-10859. U.K. AERE, Harwell, U.K.
- †Cambray, R.S., K. Playford, and N.J. Lewis. 1985. Radioactive fallout in air and rain: Results to the end of 1984. U.K. Atomic Energy Authority Rep. AERE-R-11915. U.K. AERE, Harwell, U.K.
- †Campbell, B.L. 1982. Applications of environmental caesium-137 for the determination of sedimentation rates in reservoirs and lakes and related catchment studies in developing countries. AAEC Research Establishment, Lucas Heights Research Laboratories, PMB Sutherland, 3322, N.S.W., Australia.
- †Campbell, B.L. 1983. Application of environmental caesium-137 for the determination of sedimentation rates in reservoirs and lakes and related catchment studies in developing countries. *Radioisotopes in Sediment Studies*, I.A.E.A. Tech. Doc. 298:7-30.
- Campbell, B.L., G.L. Elliott, and R.J. Loughran. 1980. Sediment dynamics in a small Hunter Valley catchment. 6:1-17. *In* Hunter environment, Univ. of Newcastle, N.S.W., Australia.
- †Campbell, B.L., G.L. Elliott, and R.J. Loughran. 1985. Nuclear fallout as an aid to measuring soil erosion. *J. Soil Conserv.* 41:86-89.
- †Campbell, B.L., G.L. Elliott, and R.J. Loughran. 1986a. Measurement of soil erosion from fallout ^{137}Cs . *Search* 17:148-149.
- †Campbell, B.L., R.J. Loughran, and G.L. Elliott. 1982. Caesium-137 as an indicator of geomorphic processes in a drainage basin system. *Aust. Geogr. Stud.* 20:49-64.
- Campbell, B.L., R.J. Loughran, and G.L. Elliott. 1988. A method for determining sediment budgets using caesium-137. *IAHS Publ.* 174:171-179.
- †Campbell, B.L., R.J. Loughran, G.L. Elliott, and D.J. Shelly. 1986b. Mapping drainage basin sediment sources using caesium-137. *Int. Assoc. of Hydrolog. Sci., Int. Symp. on Drainage Basin Sediment Delivery*, Albuquerque, NM. *IAHS Publ.* 159:437-446.
- Campbell, B.L., and M. Ross. 1980. Determination of sedimentation rate in Stephens Creek reservoir using environmental caesium-137. Newcastle (16th) Conf. of the Inst. of Australian Geographers 16:43-51.
- Carlsson, S. 1976. Cesium-137 in a dysoligotrophic lake—A radioecological field study. *LUDD/MERI 76/1005*, Univ. of Lund, Lund, Sweden.
- Carlsson, S. 1978. A model for the movement and loss of ^{137}Cs in a small watershed. *Health Phys.* 34:33-37.
- †Carrigan, P.A., Jr., and R.J. Pickering. 1967. Radioactive materials in bottom sediments of the Clinch River. Rep. ORNL-3721. Part A and B. Oak Ridge Natl. Lab., Oak Ridge, TN.
- Carpenter, R., T.M. Beasley, D. Zahnle, and B.L.K. Somayajulu. 1987. Cycling of fallout (Pu , ^{241}Am , ^{137}Cs) and natural (U , Th , ^{210}Pb) radionuclides in Washington continental slope sediments. *Geochim. Cosmochim. Acta* 51:1897-1921.
- †Carter, M.W., and A.A. Moghissi. 1977. Three decades of nuclear testing. *Health Phys.* 33:55-71.
- Cerling, T.E., and R.R. Turner. 1982. Formation of freshwater Fe-Mn coatings on gravel and behavior of ^{60}Co , ^{90}Sr , and ^{137}Cs in a small watershed. *Geochim. Cosmochim. Acta* 46:1333-1343.
- Cerrai, E., B. Schreiber, and C. Triulzi. 1967. Vertical distribution of Sr^{90} , Ce^{144} , Pm^{147} and Eu^{155} in coastal marine sediments. *Energ. Nucl. (Paris)* 14:586-592.
- Cerrai, E., and C. Triulzi. 1969. Vertical distribution of Cs^{137} , Sb^{125} , and Co^{60} in coastal marine sediments. *Energ. Nucl. (Paris)* 16:36-40.
- Chanton, J.P., C.S. Martens, and G.W. Kipphut. 1983. Lead-210 sediment geochronology in a changing coastal environment. *Geochim. Cosmochim. Acta* 47:1791-1804.
- Charles, D.F., D.R. Whitehead, D.R. Engstrom, B.D. Fry, R.A. Hites, S.A. Norton, J.S. Owen, L.A. Roll, S.C. Schindler, J.P. Smol, A.J. Uutala, J.R. White, and R.J. Wise. 1987. Paleolimnological evidence for recent acidification of Big Moose Lake, Adirondack Mountains, NY (U.S.A.). *Biogeochemistry* 3:267-296.
- Charles, M.J., and R.A. Hites. 1987. Sediments as archives of environmental pollution trends. *Adv. Chem. Ser.* 216:365-389.
- Chebotima, M.Y., and V.F. Bochenin. 1981. ^{90}Sr and ^{137}Cs in bottom sediments of a fresh water lake. (In Russian.) *Gidrobiol. Zh.* 17:82-85.
- Chenouard, L., and C. Lalou. 1969. Gamma-ray spectrometry as a tool for rapid investigation of detritic cores. *J. Sed. Petrol.* 39:1477-1483.
- Chittenden, D.M., II. 1983. Factors affecting the soluble-suspended distribution of strontium-90 and cesium-137 in Dardanelle Reservoir, Arkansas. *Environ. Sci. Technol.* 17:26-31.
- Christensen, E.R. 1982. A model for radionuclides in sediments influenced by mixing and compaction. *J. Geophys. Res.* 87:566-572.
- Christensen, E.R., and N. Chien. 1981. Fluxes of arsenic, lead, zinc, and cadmium to Green Bay and Lake Michigan sediments. *Environ. Sci. Technol.* 15:553-558.
- Christensen, E.R., and C.-K. Lo. 1986. Polychlorinated biphenyls in dated sediments of Milwaukee Harbour, Wisconsin, U.S.A. *Environ. Pollut. Ser. B* 12:217-232.
- Church, M. 1980. Records of recent geomorphological events. p. 13-29. *In* R.A. Cullingford et al. (ed.) *Timescales in geomorphology*. John Wiley & Sons, New York.
- Clifton, R.J., and E.I. Hamilton. 1982. The application of radioisotopes in the study of estuarine sedimentary processes. *Estuarine Coastal Shelf Sci.* 14:433-446.
- †Cline, J.F., and W.H. Rickard. 1972. Radioactive strontium and cesium in cultivated and abandoned field plots. *Health Phys.* 23:317-324.
- Coard, M.A., A.M. Cousen, A.H. Cuttler, H.J. Dean, J.A. Dearing, T.I. Englington, A.M. Greaves, K.P. Lacey, P.E. O'Sullivan, and D.D. Pickern. 1983. Paleolimnological studies of annually-laminated sediments in Loe Pool, Cornwall, U.K. *Hydrobiologia* 103:185-191.
- Cochran, J.K., and J.K. Osmond. 1974. Gamma spectrometry of deep-sea cores and sediment accumulation rates. *Deep Sea Res.* 21:721-737.
- †Coleman, N.T., R.J. Lewis, and D. Craig. 1963. Sorption of cesium by soils and its displacement by salt solutions. *Soil Sci. Soc. Am. Proc.* 27:290-294.
- Cooper, C.M., and J.R. McHenry. 1989. Sediment deposition and its effects on a Mississippi River oxbow lake. *Environ. Geol. Water Sci.* 13:33-37.
- Cooper, J.R., J.W. Gilliam, R.B. Daniels, and W.P. Robarge. 1987. Riparian areas as filters for agricultural sediment. *Soil Sci. Soc. Am. J.* 51:416-420.
- Cooper, J.R., J.W. Gilliam, and T.C. Jacobs. 1986. Riparian areas as a control of nonpoint pollutants. p. 166-190. *In* D.L. Correll (ed.) *Watershed research perspectives*. Smithsonian Inst. Press, Washington, DC.
- Coughtrey, P.J., and M.C. Thorne. 1983. Radionuclide distribution and transport in terrestrial and aquatic ecosystems. p. 321-424. *In* A critical review of data. Vol. 1. A.A. Balkema, Rotterdam, Netherlands.
- Cranwell, P.A. 1976. Decomposition of aquatic biota and sediment formation: Lipid components of blue-green algal species and detritus resulting from microbial attack. *Freshwater Biol.* 6:481-488.
- Cremers, A., A. Elsen, P. De Preter, and A. Maes. 1988. Quantitative analysis of radiocesium retention in soils. *Nature (London)* 335:247-249.
- Cushing, C.E., and D.G. Watson. 1974. Aquatic studies of Gable Mountain Pond. BNWL-1884, Battelle, Pacific Northwest Lab., Richland, WA.
- Cutshall, N.H., I.L. Larsen, and M.M. Nichols. 1981. Man-made radionuclides confirm rapid burial of Kepone in James River sediments. *Science (Washington, DC)* 213:440-442.
- Czuczwa, J.M., and R.A. Hites. 1984. Environmental fate of combustion-generated polychlorinated dioxins and furans. *Environ. Sci. Technol.* 18:444-450.
- †Dahlman, R.C., and S.I. Auerbach. 1968. Preliminary estimations of erosion and radiocesium redistribution in a fescue meadow. USAEC Rep. ORNL-TM-2343. Oak Ridge Natl. Lab., Oak Ridge, TN.
- †Dahlman, R.C., C.W. Francis, and T. Tamura. 1975. Radiocesium cycling in vegetation and soil. p. 462-481. *In* F.G. Howell et al. (ed.) *Mineral cycling in southeastern ecosystems*. ERDA Symp. Ser. CONF-740513. US ERDA, Washington, DC.
- Davidson, C.I., J.R. Harrington, M.J. Stephenson, M.C. Monaghan, J. Pudykiewicz, and W.R. Shell. 1987. Radioactive cesium from the Chernobyl accident in the Greenland ice sheet. *Science (Washington, DC)* 237:633-634.
- Davis, F.W. 1985. Historical changes in submerged macrophyte communities of upper Chesapeake Bay. *Ecology* 66:981-993.
- †Davis, J.J. 1963. Cesium and its relationship to potassium in ecology. p. 539-556. *In* V. Schultz and A.W. Klement, Jr. (ed.) *Radioecology*. Reinhold, New York.
- †Davis, R.B., C.T. Hess, S.A. Norton, D.W. Hanson, K.D. Hoagland, and D.S. Anderson. 1984. ^{137}Cs and ^{210}Pb dating of sediments from soft-water lakes in New England (U.S.A.) and Scandinavia, a failure of ^{137}Cs dating. *Chem. Geol.* 44:151-181.
- Davis, R.B., and S.A. Norton. 1978. Paleolimnology of six New England lakes. p. 217-308. *In* R.W. Hardy (ed.) *The impact of urbanization of New England lakes, selected technical papers*. Vol.

- III. New England Council Water Cent. Dir., Boston, MA.
- Davis, R.B., S.A. Norton, D.F. Brakke, F. Berge, and C.T. Hess. 1980. Atmospheric deposition in Norway during the last 300 years as recorded in SNSF lake sediments. Part IV. Synthesis, and comparison with New England. p. 274-275. *In* D. Drablos and A. Tøllan (ed.) Proc., int. conf. on ecological impact of acid precipitations, Sandefjord, Norway. Norway Proj. on Acid Precip.—Effects on Forest and Fish (S.N.S.F.), Norway Inst. Water Res. (N.I.V.A.), Oslo.
- Davis, R.B., S.A. Norton, C.T. Hess, and D.F. Brakke. 1983. Paleolimnological reconstruction of the effects of atmospheric deposition of acids and heavy metals on the chemistry and biology of lakes in New England and Norway. *Hydrobiologia* 103:113-123.
- Dawson, R., and E.K. Duursma. 1974. Distribution of radioisotopes between phytoplankton, sediment and sea water in a dialysis compartment system. *Neth. J. Sea Res.* 8:339-353.
- Day, L.R., and H.H. Zumpfe. 1986. Radioactivity in silt from the River Lea, England. *Environ. Pollut. Ser. B* 12:75-84.
- Dearing, J.A., J.K. Elner, and C.M. Hapley-wood. 1981. Recent sediment flux and erosional processes in a Welch upland lake-catchment based on magnetic susceptibility measurements. *Quat. Res. (NY)* 16:356-372.
- Deevey, E.S., Jr. 1988. Estimation of downward leakage from Florida lakes. *Limnol. Oceanogr.* 33:1308-1320.
- †DeJong, E., C.B.M. Begg, and R.G. Kachanoski. 1983. Estimates of soil erosion and deposition for some Saskatchewan soils. *Can. J. Soil Sci.* 63:607-617.
- DeJong, E., and R.G. Kachanoski. 1988. The importance of erosion in the carbon balance of prairie soils. *Can. J. Soil Sci.* 68:111-119.
- †DeJong, E., H. Villar, and J.R. Bettany. 1982. Preliminary investigations on the use of ^{137}Cs to estimate erosion in Saskatchewan. *Can. J. Soil Sci.* 62:673-683.
- †DeJong, E., C. Wang, and H.W. Rees. 1986. Soil redistribution on three cultivated New Brunswick hillslopes calculated from ^{137}Cs measurements, solum data and USLE. *Can. J. Soil Sci.* 66:721-730.
- DeLaune, R.D. 1986. The use of ^{13}C signature of C-3 and C-4 plants in determining past depositional environments in rapidly accreting marshes of the Mississippi River deltaic plain, Louisiana, U.S.A. *Chem. Geol.* 59:315-320.
- DeLaune, R.D., R.H. Baumann, and J.G. Gosselink. 1983. Relationships among vertical accretion, coastal submergence, and erosion in Louisiana gulf coast marsh. *J. Sed. Petrol.* 53:147-157.
- DeLaune, R.D., W.H. Patrick, Jr., and R.J. Buresh. 1978. Sedimentation rates determined by ^{137}Cs dating in a rapidly accreting salt marsh. *Nature (London)* 275:532-533.
- DeLaune, R.D., W.H. Patrick, Jr., and S.R. Pezeshki. 1987a. Foreseeable flooding and death of coastal wetland forests. *Environ. Conserv.* 14:129-133.
- DeLaune, R.D., C.N. Reddy, and W.H. Patrick, Jr. 1981. Accumulation of plant nutrients and heavy metals through sedimentation processes and accretion in a Louisiana salt marsh. *Estuar.* 4:328-334.
- DeLaune, R.D., C.J. Smith, and W.H. Patrick, Jr. 1986a. Sedimentation patterns in a Gulf Coastal backbarrier marsh: Response to increasing submergence. *Earth Surface Processes Landforms* 11:485-490.
- DeLaune, R.D., C.J. Smith, W.H. Patrick, Jr., and H.H. Roberts. 1987b. Rejuvenated marsh and bay-bottom accretion on rapidly subsiding coastal plain of U.S. Gulf Coast: A second-order effect of the emerging Atchafalaya delta. *Estuarine Coastal Shelf Sci.* 25:381-389.
- DeLaune, R.D., C.J. Smith, and M.N. Sarafyan. 1986b. Nitrogen cycling in a freshwater marsh of *Panicum hemitomon* on the deltaic plain of the Mississippi River. *J. Ecol.* 74:249-256.
- Delle Site, A., O. Ferretti, V. Marchionni, and C. Papucci. 1984. Measurements of cesium-137, plutonium-239, -240 and plutonium-238 in some coastal sediments of Tyrrhenian Sea. *Comm. Eur. Communities, (Rep.) Eur(CECED9)*, (EUR 9214, Int. Symp.), p. 233-243.
- †DeMaster, D.J., B.A. McKee, C.A. Nittrouer, D.C. Brewster, and P.E. Biscaye. 1985. Rates of sediment reworking at the Hebble site based on measurements of Th-234, Cs-137 and Pb-210. *Mar. Geol.* 66:133-148.
- Desloges, J.R. 1987. Paleohydrology of the Bella Coola river basin: An assessment of environmental reconstruction. Ph.D. thesis. Univ. of British Columbia.
- Dickman, M., H.G. Thode, S. Rao, and R. Anderson. 1988. Downcore sulphur isotope ratios and diatom inferred pH in an artificially acidified Canadian shield lake. *Environ. Pollut.* 49:265-288.
- Dickman, M.D., and H.G. Thode. 1985. The rate of lake acidification in four lakes north of Lake Superior and its relationship to downcore sulphur isotope ratios. *Water Soil Air Pollut.* 26:233-253.
- Dickson, J.H., D.A. Stewart, R. Thompson, G. Turner, M.S. Baxter, N.D. Drndarsky, and J. Rose. 1978. Palynology, paleomagnetism, and radiometric dating of Flandrian marine and freshwater sediments of Loch Lomond. *Nature (London)* 274:548-553.
- Digerfeldt, G., R.W. Battarbee, and L. Bengtsson. 1975. Report on annually laminated sediment in Lake Jarlasjon, Nacka, Stockholm. *Geol. Foeren. Stockholm Foerh.* 97:29-40.
- Dominik, J., D. Burrus, and J.-P. Vernet. 1987. Transport of the environmental radionuclides in an alpine watershed. *Earth Planet. Sci. Lett.* 84:165-180.
- Dominik, J., U. Forstner, A. Mangini, and H. Reineck. 1978. ^{210}Pb and ^{137}Cs chronology of heavy metal pollution in a sediment core from the German Bight (North Sea). *Senckenbergiana Marit* 10:213-227.
- Dominik, J., A. Mangini, and G. Muller. 1981. Determination of recent deposition rates in Lake Constance with radioisotopic methods. *Sedimentology* 28:653-677.
- Durham, R.W., and S.R. Joshi. 1980a. Recent sedimentation rates, ^{210}Pb fluxes, and particle settling velocities in Lake Huron, Laurentian Great Lakes. *Chem. Geol.* 31:53-66.
- Durham, R.W., and S.R. Joshi. 1980b. The ^{210}Pb and ^{137}Cs profiles in sediment cores from Lake Matagami and Quevillon, northwest Quebec, Canada. *Can. J. Earth Sci.* 17:1746-1750.
- Durham, R.W., and S.R. Joshi. 1981. Sedimentation rates in western Nipigon Bay, Lake Superior using the ^{210}Pb method. *Sci. Total Environ.* 22:51-59.
- Durham, R.W., and S.R. Joshi. 1984a. Dose equivalent commitments from fallout radionuclides in the open waters of the Great Lakes. 1973-1981. *Environ. Monit. Assess.* 4:405-417.
- Durham, R.W., and S.R. Joshi. 1984b. Lead-210 dating of sediments from some northern Ontario lakes. p. 75-85. *In* W.C. Mahaney (ed.) *Quaternary dating methods*. Elsevier Science Publ. B.V., Amsterdam.
- Durham, R.W., S.R. Joshi, and R.J. Allan. 1980. Radioactive dating of sediment cores from four contiguous lakes in Saskatchewan, Canada. *Sci. Total Environ.* 15:65-71.
- Durham, R.W., and B.G. Oliver. 1983. History of Lake Ontario contamination from the Niagara River by sediment radiodating and chlorinated hydrocarbon analysis. *J. Great Lakes Res.* 9:160-168.
- Duursma, E.K., and C.J. Bosch. 1970. Theoretical experimental and field studies concerning diffusion of radioisotopes in sediments and suspended particles of the sea. Part B: Methods and experiments. *Neth. J. Sea Res.* 4:395-469.
- Eadie, B.J., and J.A. Robbins. 1987. The role of particulate matter in the movement of contaminants in the Great Lakes. *Adv. Chem. Ser.* 216:319-364.
- Eakins, J.D., R.S. Cambray, K.C. Chambers, and A.E. Lally. 1984. The transfer of natural and artificial radionuclides to Brotherton water from its catchment. p. 145-164. *In* E.Y. Haworth and J.W.G. Lund (ed.) *Lake sediment and environmental history*. Univ. of Minnesota Press, Minneapolis, MN.
- Eckblad, J.W., N.L. Peterson, K. Ostlie, and A. Temte. 1977. The morphometry, benthos, and sedimentation rates of a floodplain lake in pool 9 of the upper Mississippi River. *Am. Midl. Nat.* 97:433-443.
- Edberg, N. 1980. Determination of recent sedimentation rates in lakes by ^{137}Cs . *Vatten* 36:35-44.
- Edgington, D.N., J.C. Ritchie, and J.A. Robbins. 1975. Comments on the paper, "Use of rivers to predict accumulation in sediments of radionuclides discharged from nuclear power plants," by P. Plato. *Health Phys.* 29:429-431.
- Edgington, D.N., and J.A. Robbins. 1974a. Determination of recent sedimentation rates in Lake Michigan using Pb-210 and Cs-137. p. 39-62. *In* Radiological and Environmental Research Division Annual Report, Ecology, January-December 1973. Argonne National Lab., Argonne, IL.
- Edgington, D.N., and J.A. Robbins. 1974b. The application of natural and fallout radionuclides to studies of sedimentation in Lake Michigan. p. 443-458. *In* Proc. Conf. on Nuclear Methods in Environ. Res., US ERDA Conf. 740701. US ERDA, Washington, DC.
- Edgington, D.N., and J.A. Robbins. 1975. The behavior of plutonium and other long-lived radionuclides in Lake Michigan: II. Patterns of deposition in the most recent sediments. IAEA-SM-198/40. Int. Atomic Energy Agency, Vienna.
- Edgington, D.N., and J.A. Robbins. 1976a. Patterns of deposition of natural and fallout radionuclides in the sediments of Lake Michigan and their relation to limnological processes. p. 705-729. *In* J.O. Nriagu (ed.) *Environmental biochemistry*. Vol. 2. Metals transfer and ecological mass balance. Ann Arbor Science Publ., Ann Arbor, MI.

- Edgington, D.N., and J.A. Robbins. 1976b. Patterns of deposition of radionuclides in sediments. p. 1-13. *In* Nonbiological transport and transformation of pollutants on land and water: Processes and critical data required for predictive description. National Bureau of Standards, Gaithersburg, MD.
- Edgington, D.N., and J.A. Robbins. 1976c. Records of lead deposition in Lake Michigan sediments since 1800. *Environ. Sci. Technol.* 10:266-274.
- Edgington, D.N., and J.A. Robbins. 1979. Reply to Appleby and Oldfield (1979). *Environ. Sci. Technol.* 13:480-482.
- Edgington, D.N., and J.A. Robbins. 1990. Time scales of sediment focussing in Lake Michigan as revealed by measurement of fallout Cs-137. p. 212-224. *In* M. Tilzer (ed.) Large lakes: Ecological structure and function. Science Technology, Madison, WI.
- Edgington, D.N., J.A. Robbins, and J.O. Karttunen. 1974. The distribution of Pb-210 and stable lead in Lake Michigan sediments. p. 63-76. *In* Radiological and Environmental Research Division Annual Report, Ecology. January-December 1973. Argonne National Lab., Argonne, IL.
- Efler, S.W. 1987. The impact of a chlor-alkali plant on Onondago Lake and adjoining systems. *Water Air Soil Pollut.* 33:85-115.
- Efler, S.W., and C.F. Carter. 1987. Spatial variability in selected physical characteristics and processes in Cross Lake, New York. *Water Resour. Bull.* 23:243-249.
- Efler, S.W., M.C. Rand, and T.A. Tamayo. 1979. The effect of heavy metals and other pollutants on the sediments of Onondago Lake. *Water Air Soil Pollut.* 12:117-134.
- †Elliott, G.L., B.L. Campbell, and R.J. Loughran. 1984. Correlation of erosion and erodibility assessments using Caesium-137. *J. Soil Conserv. (New South Wales)* 40:24-29.
- Elliott, G.L., R.D. Lang, and B.L. Campbell. 1983. The association of tree species, landform, soils and erosion on Narrabeen sandstone west of Putty, New South Wales. *Austr. J. Ecol.* 8:321-331.
- Ellis, J., and S. Kanamori. 1977. Water pollution studies on Lake Illawarra, III. Distribution of heavy metals in sediments. *Aust. J. Mar. Freshwater Res.* 28:485-496.
- Engstrom, D.R., E.B. Swain, and J.C. Kingston. 1985. A paleolimnological record of human disturbance from Harvey's Lake, Vermont: Geochemistry, pigments and diatoms. *Freshwater Biol.* 15:261-288.
- Erten, H.N., H.R. von Gunten, E. Rossler, and M. Sturm. 1985. Dating of sediments from Lake Zurich (Switzerland) with ²¹⁰Pb and ¹³⁷Cs. *Schweiz. Z. Hydrol.* 47:5-11.
- Essington, E.H., and E.M. Romney. 1986. Mobilization of ¹³⁷Cs during rainfall simulation studies at the Nevada Test Site. p. 35-38. *In* L.J. Lane (ed.) Proceedings of the Rainfall Simulator Workshop, Tucson, AZ. 14-15 Jan. 1985. Soc. for Range Manage., Denver, CO.
- †Evans, D.W., J.J. Alberts, and R.A. Clark III. 1983. Reversible ion-exchange fixation of cesium-137 leading to mobilization from reservoir sediments. *Geochim. Cosmochim. Acta* 47:1041-1049.
- Evans, H.E., D.C. Lasenby, and P.J. Dillon. 1986. The effect of core compression on the measurement of zinc concentrations and anthropogenic burdens in lake sediments. *Hydrobiologia* 132:185-192.
- †Eyman, L.D., and N.R. Kevern. 1975. Cesium-137 and stable cesium in a hypertrophic lake. *Health Phys.* 28:549-555.
- Farmer, J.G. 1977. The determination of sedimentation rates in Lake Ontario using the 210-Pb dating method. *Can. J. Earth Sci.* 15(3):431-437.
- Farmer, J.G. 1983. Metal pollution in marine sediment cores from the coast of Scotland. *Mar. Environ. Res.* 8:1-28.
- Farmer, J.G., D.S. Swan, and M.S. Baxter. 1980. Records and sources of metal pollutants in a dated Loch Lomond sediment core. *Sci. Total Environ.* 16:131-147.
- Favarger, P., and J. Vernet. 1979. L'isotope ¹³⁷Cs utilise comme dateur de la pollution des sediments lacustres. (In French.) *Arch. Sci.* 32:25-41.
- Feely, H.W., H.L. Volchok, E.P. Hardy, Jr., and L.E. Toonrel. 1978. Worldwide deposition of ⁹⁰Sr through 1976. *Am. Chem. Soc.* 12:808-809.
- Feijtel, T.C., R.D. DeLaune, and W.H. Patrick, Jr. 1985. Carbon flow in coastal Louisiana. *Mar. Ecol. Prog. Ser.* 24:255-260.
- Feijtel, T.C., R.D. DeLaune, and W.H. Patrick, Jr. 1988a. Biogeochemical control on metal distribution and accumulation in Louisiana sediments. *J. Environ. Qual.* 17:88-94.
- Feijtel, T.C., R.D. DeLaune, and W.H. Patrick, Jr. 1988b. Seasonal pore water dynamics in marshes of Barataria Basin, Louisiana. *Soil Sci. Soc. Am. J.* 52:59-67.
- †Fisher, J.B., W.J. Lick, P.L. McCall, and J.A. Robbins. 1980. Vertical mixing of lake sediments by tubificid Oligochaetes. *J. Geophys. Res.* 85:3997-4006.
- Flower, R.J., R.W. Battarbee, J. Natkanski, B. Rippey, and P.G. Appleby. 1988. The recent acidification of a large Scottish loch located partly within a national nature reserve and site of special scientific interest. *J. Appl. Ecol.* 25:715-724.
- Forstner, U. 1987. Sediment-associated contaminants—an overview of scientific bases for developing remedial options. *Hydrobiologia* 149:221-246.
- Foster, I.D.L., J.A. Dearing, and P.G. Appleby. 1986. Historical trends in catchment yields: A case study in reconstruction from lake-sediment records in Warwickshire, U.K. *J. Hyrol. Sci.* 31:427-443.
- Foster, I.D.L., J.A. Dearing, A. Simpson, A.D. Carter, and P.G. Appleby. 1985. Lake catchment based studies of erosion and denudation in the Merevale catchment, Warwickshire, U.K. *Earth Surface Processes Landforms* 10:45-68.
- Foster, G.R., and T.E. Hakanson. 1984. Predicted erosion and sediment delivery of fallout plutonium. *J. Environ. Qual.* 13:595-602.
- †Francis, C.W., and F.S. Brinkley. 1976. Preferential adsorption of ¹³⁷Cs to micaceous minerals in contaminated freshwater sediment. *Nature (London)* 260:511-513.
- Fredericks, D.J., V. Norris, and S.J. Perrens. 1988. Estimating erosion using cesium-137: I. Measuring caesium-137 activity in soil. *IAHS Publ.* 174:225-231.
- Fredericks, D.J., and S.J. Perrens. 1988. Estimating erosion using caesium-137: II. Estimating rates of soil loss. *IAHS Publ.* 174:233-240.
- †Fredriksson, L., B. Eriksson, B.G. Rasmuson, K. Eduarson, B. Gahne, and K. Low. 1958. Studies on soil-plant-animal interrelationships with respect to fission products, peaceful uses of atomic energy. p. 449-470. *In* Proc. of 2nd Int. Conf., Geneva, Switzerland. 1-13 September. United Nations, New York.
- †Frere, M.H., and H. Roberts, Jr. 1963. The loss of strontium-90 from small cultivated watersheds. *Soil Sci. Soc. Am. Proc.* 27:82-83.
- Gaggino, G.F., G. Parise, G. Premazzi, and A. Provini. 1984. Trends in evolution of nutrient and trace elements in sediments from 13 Italian subalpine lakes over the last 30 years. p. 135-138. *In* Proc. 3rd. Int. Symp. on the Interaction between Sediments and Water, Geneva. United Nations, New York.
- Galloway, J.N., and G.E. Likens. 1979. Atmospheric enhancement of metal deposition in Adirondack lake sediments. *Limnol. Oceanogr.* 24:427-433.
- †Garten, C.T., Jr., J.B. Gentry, J.E. Pinder III, R.R. Sharitz, and M.H. Smith. 1975. Radiocaesium dynamics in a contaminated flood plain ecosystem in the southeastern United States. p. 331-347. *In* Impacts of nuclear releases into the aquatic environment. Int. Atomic Energy Agency, Vienna.
- Giani, L., H. Gebhardt, W. Gusy, and H. Helmers. 1987. Verhalten einiger radioaktiver nuklide (freigesetzt durch den reaktorunfall in Tschernobyl) in typischen Boden Norddeutschlands. *Z. Pflanzenernaehr. Bodenkd.* 150:103-107.
- Gilbert, R., and J.R. Desloges. 1988. Sediments of ice-dammed, self-draining Ape Lake, British Columbia. *Can. J. Earth Sci.* 24:1735-1747.
- Gilbey, J., S. Bradley, and D.E. Walling. 1987. The deposition of cesium 137 on grassland at a site in SW England following the Chernobyl accident. *Grass Forage Sci.* 42:439-442.
- Gill, A.C., J.R. McHenry, and J.C. Ritchie. 1976. Efficiency of nitrogen, carbon, and phosphorus retention by small agricultural reservoirs. *J. Environ. Qual.* 5:310-315.
- Goreau, T.J. 1977. Quantitative effects of sediment mixing on stratigraphy and biogeochemistry: A signal theory approach. *Nature (London)* 265:525-526.
- †Graham, E.R. 1963. Factors affecting Sr-85 and I-131 removal by runoff water. *Water Sewage Works* 110:407-410.
- Gregorich, E.G., and D.W. Anderson. 1985. Effects of cultivation and erosion on soils of four toposequences in the Canadian prairies. *Geoderma* 36:343-354.
- Griffiths, M. 1978. Specific blue-green algal carotenoids in sediments of Esthwaite Water. *Limnol. Oceanogr.* 23:777-784.
- Grimes, J.A., and S.R. Rushforth. 1983. Diatoms of surface sediments of Utah Lake, Utah, U.S.A. *Hydrobiologia* 99:161-174.
- Grzybowska, D., and G.J. Soszka. 1978. ⁹⁰Sr and ¹³⁷Cs contamination level in Zarnowiec Lake. *Bulletin de L'Academie Polonaise des Sciences Serie des Sciences Biologiques Cl. II* 26:751-757.
- Gschwend, P.M., and R.A. Hites. 1981. Fluxes of polycyclic aromatic hydrocarbons to marine and lacustrine sediments in the north-eastern United States. *Geochim. Cosmochim. Acta* 45:2359-2367.
- Haghiri, F. 1970. Fate of strontium-90 in soil, plants, and water. *Ohio Rep.* 55:74-77. Ohio Agric. Res. and Dev. Center, Wooster, OH.
- Hairr, L. 1974. An investigation of factors influencing radiocesium cycling in estuarine sediments of the Hudson River. Ph.D. thesis. New York Univ. (DAI no. 76:19605).
- Hakanson, L. 1980. An ecological risk index for aquatic pollution

- control. A sedimentological approach. *Water Res.* 14:975-1001.
- †Hakanson, L., and A. Kallstrom. 1978. An equation of state for biologically active lake sediments and its implications for interpretations of sediment data. *Sedimentology* 25:205-226.
- Hakkinen, U., and E. Lakanen. 1968. Strontium-90 and caesium-137 in some Finnish soil profiles. *Ann. Agric. Fenn.* 7:123-126.
- Hamilton, E.I., and K.R. Clarke. 1984. The recent sedimentation history of the Esk estuary, Cumbria, U.K.: The application of radiochronology. *Sci. Total Environ.* 35:325-386.
- Hardy, E.P., H.D. Livingston, J.C. Burke, and H.L. Volchok. 1978. Time pattern of off-site plutonium contamination from Rocky Flats plants by lake sediment analyses. USDOE Rep. EML-342. p. I-123-146. EML, New York.
- Harris, C. 1977. Cs-137 as a tool for measuring erosion in the Lake Thunderbird watershed: A feasibility study. Unpublished senior study, Univ. of Oklahoma, Norman, OK.
- Haselwandter, K. 1977. Accumulation of the radioactive nuclide ^{137}Cs in fruitbodies of basidiomycetes. *Health Phys.* 34:713-715.
- Hatton, R.S., R.D. DeLaune, and W.H. Patrick, Jr. 1983. Sedimentation, accretion, and subsidence in marshes of Barataria Basin, Louisiana. *Limnol. Oceanogr.* 28:494-502.
- Haworth, E.Y. 1976. The changes in the composition of diatom assemblages found in the surface sediments of Blelham Tarn in the English Lake District during 1973. *Ann. Bot.* 40:1195-1205.
- Haworth, E.Y. 1980. Comparison of continuous phytoplankton records with the diatom stratigraphy in recent sediments of Blelham Tarn. *Limnol. Oceanogr.* 25:1093-1103.
- Haworth, E.Y. 1984. Stratigraphic changes in algal remains (diatoms and chrysophytes) in recent sediments of Blelham Tarn, English Lake District. p. 165-190. In E.Y. Haworth and J.W.G. Lund (ed.) *Lake sediments and environmental history*. Univ. of Minnesota Press, Minneapolis, MN.
- †Health and Safety Laboratory Fallout Program. 1972. USAEC, HASL-258, A-40-41, A-93-94. Health and Safety Lab., New York.
- Hebel, S.J., and D.C. Wilkin. 1982. Agricultural land management strategies for sediment reduction. *Landscape Architecture*, September, p. 91-95.
- Heit, M., and K.M. Miller. 1987. Cesium-137 sediment depth profiles and inventories in Adirondack lake sediments. *Biogeochemistry* 3:243-265.
- Heit, M., K.M. Miller, P. Krey, D. Bogen, and H. Feely. 1984. Other radionuclides: Plutonium with some comments about ^{137}Cs chronologies in freshwater sediments. p. 34-77. In *Workshop on Paleolimnological Studies of the History and Effects of Acidic Precipitation*, Rockland, ME. 23-25 May. USEPA, Washington, D.C.
- Heit, M., Y. Tan, C. Klusek, and J.C. Burke. 1981. Anthropogenic trace elements and polycyclic aromatic hydrocarbon levels in sediment cores from two lakes in the Adirondack acid lake region. *Water Air Soil Pollut.* 15:441-464.
- Heit, M., Y.L. Tan, C.S. Klusek, H.L. Volchok, and J.C. Burke. 1980. The origin and deposition history of trace elements and polynuclear aromatic hydrocarbons in two remote lakes in the Adirondack acid rain region. USDOE Rep. EML-381, p. I-75 to I-127. EML, New York.
- Helton, J.C., A.B. Muller, and A. Bayer. 1985. Contamination of surface-water bodies after reactor accidents by the erosion of atmospherically deposited radionuclides. *Health Phys.* 48:757-771.
- Hetherington, J.A., and D.F. Jefferies. 1974. The distribution of some fission product radionuclides in sea and estuarine sediments. *Neth. J. Sea Res.* 8:319-338.
- Hilton, J., J.P. Lishman, and P.V. Allen. 1986. The dominant processes of sediment distribution and focusing in a small, eutrophic, monomictic lake. *Limnol. Oceanogr.* 31:125-133.
- Mierschberg, D.J., and J.R. Schubel. 1979. Recent geochemical history of flood deposits in the northern Chesapeake Bay. *Estuarine Coastal Mar. Sci.* 9:771-784.
- Hubbard, J.E., and W.D. Striffler. 1973. Cesium-137 in a mountain stream channel. *Water Resour. Res.* 9:1440-1442.
- Hutchison-Benson, E., J. Svoboda, and H.W. Taylor. 1985. The latitudinal inventory of ^{137}Cs in vegetation and topsoil in northern Canada, 1980. *Can. J. Bot.* 63:784-791.
- Inbar, M. 1982. Measurement of fluvial sediment transport compared with lacustrine sedimentation rates: The flow of the River Jordan into Lake Kinneret. *J. Hydrol. Sci.* 4:439-449.
- Irlweck, K. 1985. Depth distribution of $^{137}\text{Cesium}$, $^{90}\text{Strontium}$ and $^{210}\text{Lead}$ in sediments of Lake Mondsee, Austria. *J. Radioanal. Nucl. Chem. Lett.* 93:115-124.
- Irlweck, K., and D.L. Danielopol. 1985. Caesium-137 and lead-210 dating of recent sediments from Mondsee (Austria). *Hydrobiologia* 128:175-185.
- Jaakkola, T., K. Tolonen, P. Huttunen, and S. Leskinen. 1983. The use of fallout ^{137}Cs and $^{239,240}\text{Pu}$ for dating of lake sediments. *Hydrobiologia* 103:15-19.
- Jasinski, T., and J.A. Robbins. 1988. Radiotracers in Polish lake sediments. *EOS Trans.* 69:1119.
- Jaworowski, Z. 1966. Temporal and geographical distribution of radium D (Lead-210). *Nature (London)* 212:886-889.
- Jenne, E.A., and J.S. Wahlberg. 1968. Role of certain stream-sediment components in radioion sorption. U.S. Geol. Surv. Prof. Pap. 433-F.
- Johnson, T.C. 1984. Sedimentation in large lakes. *Ann. Rev. Earth Planet. Sci.* 12:179-204.
- Johnston, C.A., G.D. Bubenzer, G.B. Lee, F.W. Madison, and J.R. McHenry. 1984. Nutrient trapping by sediment deposition in a seasonally flooded lakeside wetland. *J. Environ. Qual.* 13:283-290.
- Johnston, S.E. 1981. A comparison of dating methods in laminated lake sediments in Maine. M.S. thesis. Univ. of Maine, Orono.
- Johnston, S.E., S.A. Norton, C.T. Hess, R.B. Davis, and R.S. Anderson. 1982. Chronology of atmospheric deposition of acids and metals in New England, based on the record in lake sediments. p. 177-187. In L.H. Keith (ed.) *Energy and environmental chemistry. Vol. 2. Acid rain*. Ann Arbor Science Publ., Ann Arbor, MI.
- Jones, D.G., J.M. Miller, and P.D. Roberts. 1984. The distribution of ^{137}Cs in surface intertidal sediments from Solway Firth. *Mar. Pollut. Bull.* 15:187-194.
- Joshi, S.R. 1985. Recent sedimentation rates and ^{210}Pb fluxes in Georgian Bay and Lake Huron. *Sci. Total Environ.* 41:219-233.
- Joshi, S.R. 1988. West Valley-derived radionuclides in the Niagara River area of Lake Ontario. *Water Air Soil Pollut.* 37:111-120.
- Joshi, S.R., and R.W. Durham. 1976. Determination of ^{210}Pb , ^{226}Ra and ^{137}Cs in sediments. *Chem. Geol.* 18:155-160.
- Joshi, S.R., and M.E. Fox. 1985. The ^{210}Pb and ^{137}Cs profiles in sediment cores from Bay of Quinte, Lake Ontario. *J. Radioanal. Nucl. Chem. Articles* 90:207-215.
- Joshi, S.R., and A. Mudroch. 1988. Direct determination of geochronologically useful radionuclides in sediments by low-energy photon analysis. *Nucl. Instrum. Methods Phys. Res.* 263:529-536.
- Juang, T.C. 1988. Tracing soil erosion sources of Teh-chi reservoir watershed by radiochemical and geochemical methods. *Soil Fert. Taiwan* 1988:17-24.
- Juzdan, Z. 1988. Worldwide deposition of ^{90}Sr through 1985. USDOE Rep. EML-515. EML, New York.
- †Kachanoski, R.G. 1987. Comparison of measured soil 137-cesium losses and erosion rates. *Can. J. Soil Sci.* 67:199-203.
- †Kachanoski, R.G., and E. DeJong. 1984. Predicting the temporal relationship between soil cesium-137 and erosion rate. *J. Environ. Qual.* 13:301-304.
- †Kaciszczenko, J., and Z. Banasik. 1981. An effect of bioturbation on the results of the ^{137}Cs dating technique used for lacustrine sediments. *Ekol. Pol.* 29:615-623.
- Kadlec, R.H., and J.A. Robbins. 1984. Sedimentation and sediment accretion in Michigan coastal wetlands. *Chem. Geol.* 44:119-150.
- Kamiyama, K., S. Okuda, and M. Koyama. 1982. Vertical distribution of ^{137}Cs and its accumulation rate in lake sediments. *Jpn. J. Limnol.* 43:35-38.
- Kansanen, P.H. 1985. Assessment of pollution history from recent sediments in Lake Vanajavesi, southern Finland. II. Changes in the Chironomidae, Chaoboridae and Ceratopogonidae (Diptera) fauna. *Ann. Zool. Fenn.* 22:57-90.
- Kansanen, P.H., and T. Jaakkola. 1985. Assessment of pollution history from recent sediments in Lake Vanajavesi, southern Finland. I. Selection of representative profiles, their dating and chemostratigraphy. *Ann. Zool. Fenn.* 22:13-55.
- Kawabata, T. 1967. Studies on the sorption and release of radionuclides by river sediments. *J. Radiat. Res.* 8:20-31.
- Kedhi, K. 1989. Radiotracer study of deposition caused by suspended sediments of river Seman. *J. Radioanal. Nucl. Chem. Lett.* 135:395-402.
- Keilty, T.J., D.S. White, and P.F. Landrum. 1988. Sublethal responses to endrin in sediment by *Stylodrilus heringianus* (Lumbriculidae) as measured by a $^{137}\text{cesium}$ marker layer technique. *Aquat. Toxicol.* 13:251-270.
- Kemp, A.L.W., T.W. Anderson, R.L. Thomas, and A. Mudrochova. 1974. Sedimentation rates and recent sediment history of Lakes Ontario, Erie and Huron. *J. Sed. Petrol.* 44:207-218.
- Kiss, J.J., E. DeJong, and L.W. Martz. 1988. The distribution of fallout Cesium-137 in southern Saskatchewan, Canada. *J. Environ. Qual.* 17:445-452.
- †Kiss, J.J., E. DeJong, and H.P.W. Rostad. 1986. An assessment of soil erosion in west-central Saskatchewan using cesium-137. *Can. J. Soil Sci.* 66:591-600.
- Klechkovskii, V.M., G.G. Polikarpov, and R.M. Aleksakhin (ed.). 1972. Behavior of strontium-90 and caesium-137 in soils. p. 78-102. In *Radioecology*. John Wiley & Sons, New York.
- Klusek, C.S. 1987. Strontium-90 in food and bone from fallout. *J. Environ. Qual.* 16:195-199.

- Kodaira, K., M. Kato, and M. Ishikawa. 1973. Radiochemical analysis of ^{210}Pb , as well as distribution of ^{210}Pb , ^{90}Sr , and ^{137}Cs in soil profiles. *Radioisotopes* 22(7):331-336.
- Koh, M., and J.R. McHenry. 1982. Determining sediment rate using cesium-137 fallout at Larto Lake. *J. Korean Soc. Soil Sci. Fert.* 15:207-212.
- Koide, M., K.W. Bruland, and E.D. Goldberg. 1973. Th-228/Th-232 and Pb-210 geochronologies in marine and lake sediments. *Geochim. Cosmochim. Acta* 35:1171-1187.
- Komarneni, S. 1978. Cesium sorption and desorption behavior of kaolinites. *Soil Sci. Soc. Am. J.* 42:531-532.
- Korschgen, C.E., G.A. Jackson, L.F. Muessig, and D.C. Southworth. 1987. Sedimentation in Lake Onalaska, Navigation Pool 7, Upper Mississippi River, since impoundment. *Water Resour. Bull.* 23:221-226.
- Krey, P.W., E.P. Hardy, and M. Heit. 1980. Nevada test site fallout in the area of Enterprise, Utah. USDOE Rep. EML-372. EML, New York.
- †Krezoski, J.R., S.C. Mozley, and J.A. Robbins. 1978. Influence of benthic macroinvertebrates on mixing of profundal sediments in southeastern Lake Huron. *Limnol. Oceanogr.* 23:1011-1016.
- †Krezoski, J.R., and J.A. Robbins. 1985. Vertical distribution of feeding and particle-selective transport of ^{137}Cs in lake sediments by lumbricid oligochaetes. *J. Geophys. Res.* 90:11999-12006.
- Krezoski, J.R., J.A. Robbins, and D.S. White. 1984. Dual radiotracer measurement of zoobenthos-mediated solute and particle transport in freshwater sediments. *J. Geophys. Res.* 89:7937-7947.
- Krishnaswami, S., K. Benninger, R.C. Aller, and K.L. Von Damm. 1980. Atmospherically-derived radionuclides as tracers of sediment mixing and accumulation in near-shore marine and lake sediments: Evidence from ^7Be , ^{210}Pb and $^{239,240}\text{Pu}$. *Earth Planet. Sci. Lett.* 47:307-318.
- Krishnaswami, S., and D. Lal. 1978. Radionuclide limnology. p. 153-177. *In* A. Lerman (ed.) *Lakes; chemistry, geology, physics*. Springer-Verlag, New York.
- †Krishnaswami, S., D. Lal, J.M. Martin, and M. Meybeck. 1971. Geochronology of lake sediments. *Earth Planet. Sci. Lett.* 11:407-414.
- Kudo, A., and E.F. Gloyna. 1971. Transport of ^{137}Cs -II: Interaction with bed sediments. *Water Res.* 5:71-79.
- †Lance, J.C., S.C. McIntyre, D.G. DeCoursey, P.G. Sharpe, S.S. Rousseva, and B.L. Campbell. 1986a. Using cesium-137 to estimate erosion rates on rangeland. p. 79-89. *In* *Erosion control, protecting our future*. Int. Erosion Control Assoc., Steamboat Springs, CO.
- †Lance, S.C. McIntyre, R.R. Lowrance, B.L. Campbell, and R.L. Miller. 1986b. Cesium-137 measures erosion rate and sediment movement. p. 4-1 to 4-9. *In* 4th Federal Interagency Sedimentation Conference, Las Vegas, NV. 24-27 March. Water Resources Council, Washington, DC.
- †Lance, J.C., S.C. McIntyre, J.W. Naney, and S.S. Rousseva. 1986c. Measuring sediment movement at low erosion rates using cesium-137. *Soil Sci. Soc. Am. J.* 50:1303-1309.
- †Larsen, R.J. 1985. Worldwide deposition of Sr-90 through 1983. USDOE Rep. EML-444. EML, New York.
- Lehman, J.T. 1975. Reconstructing the rate of accumulation of lake sediment: The effect of sediment focusing. *Quat. Res. (NY)* 5:541-550.
- Lentsch, J.W. 1974. The fate of gamma-emitting radionuclides released into the Hudson River Estuary and an evaluation of their environmental significance. Ph.D. thesis. New York Univ., New York.
- Leonard, E.M. 1986. Varve studies at Hector Lake, Alberta, Canada, and the relationship between glacial activity and sedimentation. *Quat. Res. (NY)* 25:199-214.
- †Lerman, A., and T.A. Lietzke. 1975. Uptake and migration of tracers in lake sediments. *Limnol. Oceanogr.* 20:497-510.
- Lewin, J. 1977. Floodplain geomorphology. *Prog. Phys. Geog.* 1:408-214.
- Li, Y.-H., L. Burkhardt, M. Bucholtz, P. O'Hara, and P.H. Santschi. 1984. Partition of radiotracers between suspended particles and sea water. *Geochim. Cosmochim. Acta* 48:2011-2020.
- Livens, F.R., and P.J. Loveland. 1988. The influence of soil properties on the environmental mobility of caesium in Cumbria. *Soil Use Manage.* 4:69-75.
- Livingston, H.D. 1984. The preservation of algal remains in recent lake sediment. p. 191-202. *In* E.Y. Haworth and J.W.G. Lund (ed.) *Lake sediment and environmental history*. Univ. of Minnesota Press, Minneapolis, MN.
- Livingston, H.D., and V.T. Bowen. 1979. Pu and ^{137}Cs in coastal sediments. *Earth Planet. Sci. Lett.* 43:29-45.
- Livingstone, H.D., and R.S. Cambray. 1978. Confirmation of ^{137}Cs dating by algal stratigraphy in Rostherne Mere. *Nature (London)* 276:259-261.
- †Lomenick, T.F., and T. Tamura. 1965. Naturally occurring fixation of cesium-137 on sediments of lacustrine origin. *Soil Sci. Soc. Am. Proc.* 29:383-386.
- †Longmore, M.E. 1982. The caesium-137 dating technique and associated applications in Australia—a review. p. 310-321. *In* W. Ambrose and P. Duerden (ed.) *Archaeometry: An Australasian perspective*. Australian National Univ. Press, Canberra, Australia.
- Longmore, M.E., J.G. Luly, and B.M. O'Leary. 1986a. Cesium-137 redistribution in the sediments of the Playa, Lake Tyrrell, northwestern Victoria. II. Patterns of cesium-137 and pollen redistribution. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 54:197-218.
- †Longmore (McCallan), M.E., B.M. O'Leary, and C.W. Rose. 1983a. Cesium-137 profiles in the sediment of a partial-meromictic lake on Great Sandy Island (Fraser Island), Queensland, Australia. *Hydrobiologia* 103:21-27.
- †Longmore (McCallan), M.E., B.M. O'Leary, C.W. Rose, and A.L. Chandra. 1983b. Mapping soil erosion and accumulation with the fallout isotope caesium-137. *Aust. J. Soil Res.* 21:373-385.
- Longmore, M.E., T. Torgersen, B.M. O'Leary, and J.G. Luly. 1986b. Cesium-137 redistribution in the sediment of the Playa, Lake Tyrrell, northwestern Victoria. I. Stratigraphy and Cesium-137 mobility in the upper sediments. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 54:181-195.
- Loughran, R.J., and B.L. Campbell. 1983. The determination of sedimentation depth by caesium-137. *Search* 14:157-158.
- Loughran, R.J., B.L. Campbell, and G.L. Elliott. 1980. Sediment dynamics in Maluna Creek catchment at Pokolbin, N.S.W. *Papers of the Newcastle (16th) Conference of the Institute of Australian Geographers* 16:245-254.
- †Loughran, R.J., B.L. Campbell, and G.L. Elliott. 1981. Sediment erosion, storage and transport in a small steep drainage basin at Pokolbin, N.S.W., Australia. p. 252-268. *In* Proc. Christchurch Symp., Christchurch, New Zealand. 25-31 January. IAHS Publ. 132. IAHS Press, Wallingford, U.K.
- †Loughran, R.J., B.L. Campbell, and G.L. Elliott. 1982. The identification and quantification of sediment sources using ^{137}Cs . p. 361-369. *In* Proc. Exeter Symp., Exeter, U.K. 19-30 July. IAHS Publ. 137. IAHS Press, Wallingford, U.K.
- †Loughran, R.J., B.L. Campbell, and G.L. Elliott. 1986a. A nuclear technique measures soil erosion. *Nucl. Spect.* 2:2-4.
- †Loughran, R.J., B.L. Campbell, and G.L. Elliott. 1986b. The use of tracer caesium-137 for studying sediment movement in a drainage basin. p. 151-160. *In* E. Frankel et al. (ed.) *Recent sediments in eastern Australia—marine through terrestrial*. Geological Soc. of Australia, Univ. of Sydney, Australia.
- †Loughran, R.J., B.L. Campbell, and G.L. Elliott. 1986c. Sediment dynamics in a partially cultivated catchment in New South Wales, Australia. *J. Hydrol.* 83:285-297.
- †Loughran, R.J., B.L. Campbell, A.T. Pilgrim, and A.J. Conacher. 1987a. Caesium-137 in soils in relation to the nine unit landsurface model in a semi-arid environment in Western Australia. p. 398-406. *In* A. Conacher (ed.) *Readings in Australian geography*. Proc. of the 21st Inst. of Australian Geographers' Conf., Perth, Australia. 10-18 May 1986. Univ. of Western Australia, Perth.
- †Loughran, R.J., B.L. Campbell, and D.E. Walling. 1987b. Soil erosion and sedimentation indicated by caesium 137: Jackmoor Brooke catchment Devon, U.K. *Catena* 14:201-212.
- Loughran, R.J., B.L. Campbell, D.E. Walling, and G.L. Elliott. 1985. Caesium redistribution and sediment movement in two catchments in England and Australia, First Int. Conf. on Geomorphology, Manchester, U.K. (Poster Report.)
- Loughran, R.J., G.L. Elliott, B.L. Campbell, and D.J. Shelly. 1988. Estimation of soil erosion from caesium-137 measurements in a small cultivated catchment in Australia. *Appl. Radiat. Isot.* 39:1153-1157.
- †Lowrance, R., S. McIntyre, and C. Lance. 1988. Sediment deposition in a field/forest system estimated using cesium-137 activity. *J. Soil Water Conserv.* 43:195-199.
- Ludlam, S.D. 1984. Fayetteville Green Lake, New York, U.S.A. VII. Varve chronology and sediment focusing. *Chem. Geol.* 44:85-100.
- Lund, J.W.G. 1984. Winifred Tutin—a personal note. p. 1-10. *In* E.Y. Haworth and J.W.G. Lund (ed.) *Lake sediment and environmental history*. Univ. of Minnesota Press, Minneapolis, MN.
- Lynch, D.R., and C.B. Officer. 1984. Nonlinear parameter estimation for sediment cores. *Chem. Geol.* 44:203-225.
- Lyons, W.B., P.B. Armstrong, and H.E. Gaudette. 1983. Trace metal concentrations and fluxes in Bermuda sediments. *Mar. Pollut. Bull.* 14:65-68.
- Mackenzie, A.B., M.S. Baxter, I.G. McKinley, D.S. Swan, and W. Jack. 1979. The determination of ^{134}Cs , ^{137}Cs , ^{210}Pb , ^{226}Ra , and ^{228}Ra concentrations in nearshore marine sediments and seawater. *J. Radioanal. Chem.* 48:29-47.
- Mahara, Y., A. Kudo, T. Kauri, D.C. Santry, and S. Miyahara. 1988. Mobile Pu in reservoir sediments of Nagasaki, Japan. *Health*

- Phys. 54:107-111.
- Makhon'ko, K.P., Ts.I. Bobovnikova, A.S. Avramenko, A.V. Dibtseva, and A.A. Volokitin. 1975. Vertical distribution of Sr^{90} and Cs^{137} in bottom deposits of lakes and rivers. *Ekologiya* 6:90-93.
- Martin, D.B. 1985. Accumulation of sediment, nutrients, and cesium-137 in prairie potholes in cultivated and noncultivated watersheds. p. 274-275. *In* Perspectives in nonpoint source pollution. USEPA, Washington, DC.
- Martin, D.B., and W.A. Hartman. 1987. The effect of cultivation on sediment composition and deposition in prairie pothole wetlands. *Water Air Soil Pollut.* 34:45-53.
- Martin, J. 1985. The Pavin Crater Lake. p. 169-188. *In* W. Stumm (ed.) Chemical processes in lakes. John Wiley & Sons, New York.
- Martz, L.W., and E. DeJong. 1985. The relationship between land surface morphology and soil erosion and deposition in a small Saskatchewan basin. *Proc. of the Seventh Hydrotechnical Conf., Canadian Soc. for Civil Eng. Annual Conf.* 1985:1-19.
- †Martz, L.W., and E. DeJong. 1987. Using cesium-137 to assess the variability of net soil erosion and its association with topography in a Canadian prairie landscape. *Catena* 14:439-451.
- Mathewes, R.W., and J.M. D'Auria. 1982. Historic changes in an urban watershed determined by pollen and geochemical analyses of lake sediment. *Can. J. Earth Sci.* 19:2114-2125.
- Matisoff, G., and J.A. Robbins. 1987. A model for biological mixing of sediments. *J. Geol. Educ.* 35:144-149.
- Matschullat, V.J., H. Heinrichs, J. Schneider, and M. Sturm. 1987. Heavy metal contents in lake sediments of the western Harz mountains (FRG) (In German.) *Chem. Erde* 47:181-194.
- Maule, C.P., and M.J. Dudas. 1989. Preliminary identification of soil separates associated with fallout ^{137}Cs . *Can. J. Soil Sci.* 69:171-175.
- McCabe, W.J., and W.J. Mason. 1982. A Falls Dam, Otago, sedimentation rate measured by caesium-137 profile. *INS-R-309*. Inst. of Nucl. Sci., Lower Hutt, New Zealand.
- McCall, P.L., J.A. Robbins, and G. Matisoff. 1984. ^{137}Cs and ^{210}Pb transport and geochronologies in urbanized reservoirs with rapidly increasing sedimentation rates. *Chem. Geol.* 44:33-65.
- †McCallan, M.E., B.M. O'Leary, and C.W. Rose. 1980. Redistribution of caesium-137 by erosion and deposition on an Australian soil. *Aust. J. Soil Res.* 18:119-128.
- McCallan, M.E., and C.W. Rose. 1977. The construction of a geochronology for alluvial deposits on the Condamine Plain and the estimation of the areal variation in erosion intensity in the Upper Condamine drainage basin. *Tech. Rep. 2/77*. School of Australian Environ. Studies, Griffith Univ.
- McHenry, J.R. 1968. The use of tracer technique in soil erosion research. *Trans. Am. Soc. Agric. Eng.* 11:619-625.
- McHenry, J.R. 1974. Reservoir sedimentation. *Water Resour. Bull.* 10:329-337.
- †McHenry, J.R., and G.B. Bubenzer. 1985. Field erosion estimated from ^{137}Cs activity measurements. *Trans. Am. Soc. Agric. Eng.* 28:480-483.
- McHenry, J.R., C.M. Cooper, and J.C. Ritchie. 1982. Sedimentation in Wolf Lake, Lower Yazoo River Basin, Mississippi. *J. Freshwater Ecol.* 1:547-558.
- McHenry, J.R., and S.C. McIntyre. 1984. Recent sedimentation rates in two North American impoundments with watersheds predominantly in cropland. p. 205-211. *In* R.J. Loughran (ed.) Drainage basin erosion and sedimentation, conference papers. Univ. of Newcastle, and Soil Conserv. Serv., N.S.W., Australia.
- †McHenry, J.R., and J.C. Ritchie. 1975. Redistribution of cesium-137 in southeastern watersheds. p. 452-461. *In* F.G. Howell et al. (ed.) Mineral cycling in southeastern ecosystems. ERDA, Symp. Series, CONF-740513. US ERDA, Washington, DC.
- †McHenry, J.R., and J.C. Ritchie. 1977a. Estimating field erosion losses from fallout cesium-137 measurements. *IAHS-AISH Publ.* 122. IAHS Press, Wallingford, U.K., p. 26-33.
- McHenry, J.R., and J.C. Ritchie. 1977b. Physical and chemical parameters affecting transport of Cs^{137} in arid watersheds. *Water Resour. Res.* 13:923-927.
- McHenry, J.R., and J.C. Ritchie. 1980. Dating recent sediments in impoundments. p. 1279-1289. *In* Proc. symp. on surface water impoundments, Minneapolis, MN. 2-5 June. Am. Soc. Civil Eng., New York.
- †McHenry, J.R., J.C. Ritchie, and G.D. Bubenzer. 1978. Redistribution of caesium-137 due to erosional processes in a Wisconsin watershed. p. 495-503. *In* D.C. Adriano and I.L. Brisbin, Jr. (ed.) Environmental Chemistry and Cycling Processes. DOE CONF-760429. US DOE, Washington, DC.
- McHenry, J.R., J.C. Ritchie, and C.M. Cooper. 1980. Rates of recent sedimentation in Lake Pepin. *Water Resour. Bull.* 16:1049-1056.
- †McHenry, J.R., J.C. Ritchie, C.M. Cooper, and J. Verdon. 1984. Recent rates of sedimentation in the Mississippi River. p. 99-117. *In* J.G. Wiener et al. (ed.) Contaminants in the Upper Mississippi River. Butterworth Publ., Boston, MA.
- †McHenry, J.R., J.C. Ritchie, and A.C. Gill. 1973. Accumulation of fallout cesium-137 in soils and sediments in selected watersheds. *Water Resour. Res.* 9:676-686.
- McHenry, J.R., J.C. Ritchie, and J. May. 1975. Recent sedimentation rates in the lower Mississippi River Valley: Lake Verret-Lake Palourde, Louisiana. p. 13-23. *In* Proc. Miss. Water Resour. Conf., Jackson, MS. 12 April. Mississippi Water Resour. Inst., Mississippi State, MS.
- McHenry, J.R., J.C. Ritchie, and J. Verdon. 1976. Sedimentation rates in the Upper Mississippi River. p. 1339-1349. *In* Symposium on inland waterways for navigation, flood control and water diversions. Vol. 2. Am. Soc. Civil Eng., New York.
- McHenry, J.R., F.R. Schiebe, J.C. Ritchie, and C.M. Cooper. 1984. Deposited sediments in Lake Chicot. p. 18-47. *In* J.F. Nix and F.R. Schiebe (ed.) Limnological studies of Lake Chicot, Arkansas. Quachita Baptist College, Arkadelphia, AK.
- †McIntyre, S.C., J.C. Lance, B.L. Campbell, and R.L. Miller. 1987. Using cesium-137 to estimate soil erosion on a clearcut hillside. *J. Soil Water Conserv.* 42:117-120.
- McIntyre, S.C., J.W. Naney, and W.R. Berkas. 1988. Measurement of recent sediment deposition in Lake Taneycomo using cesium-137. Abstracts of Papers for the 1988 Annual Meeting of the American Society of Limnology and Oceanography, Boulder, CO.
- McIntyre, S.C., J.W. Naney, and J.R. McHenry. 1989. Quantification of lake sedimentation rates utilizing radioisotopes present in the environment. p. 1-5. *In* Offsite assessment. EPA Publ. 440/5-89-001. USEPA, Washington, DC.
- †Menzel, R.G. 1960. Transport of strontium-90 in runoff. *Science* (Washington, DC) 131:499-500.
- †Menzel, R.G., P. Jung, K. Ryu, and K. Um. 1987. Estimating soil erosion losses in Korea with fallout cesium-137. *J. Appl. Radiat. Isot.* 38:451-454.
- Merefield, J.R. 1980. Caesium in the up-estuary transport of sediments. *Mar. Geol.* 39:45-55.
- Merefield, J.R. 1986. Alkali metals in the permo-triassic as geochemical indicators of surficial processes. *Chem. Geol.* 56:143-158.
- Meriwether, J.R., J.N. Beck, D.F. Keeley, M.P. Langley, R.H. Thompson, and J.C. Young. 1988. Radionuclides in Louisiana Soils. *J. Environ. Qual.* 17:562-568.
- Miller, K.M., and M. Heit. 1986. A time resolution methodology for assessing the quality of lake sediment cores that are dated by ^{137}Cs . *Limnol. Oceanogr.* 31:1292-1300.
- †Mitchell, J.K., G.D. Bubenzer, J.R. McHenry, and J.C. Ritchie. 1980. Soil loss estimation from fallout cesium-137 measurements. p. 393-401. *In* M. DeBoodt and D. Grabriels (ed.) Assessment of erosion. John Wiley & Sons, New York.
- Mitchell, J.K., S. Mostaghimi, D.S. Freeny, and J.R. McHenry. 1983. Sediment deposition estimation from cesium-137 measurements. *AWRA, Water Resour. Bull.* 19:549-555.
- Mohamad, D.B. 1982. Application of Cs^{137} techniques to problems of sediment redistribution in Sungai Lui representative basin, Selangor, Malaysia: Part I. IAEA Regional Coop. Agreement Rep., MAL 2623/RI/AG. Int. Atomic Energy Agency, Vienna.
- Moore, P.D. 1974. Lead accretion in alluvial deposits. *Nature* (London) 250:288-289.
- Muller, R.M., D.G. Sprugel, and B. Kohn. 1978. Erosional transport and deposition of plutonium and cesium in two small midwestern watersheds. *J. Environ. Qual.* 7:171-174.
- Nagaya, Y., and M. Saiki. 1967. Accumulation of radionuclides in coastal sediments of Japan (I) Fallout radionuclides in some coastal sediments in 1964-1965. *J. Radiat. Res.* 8:37-43.
- Nakamura, K., and Y. Nagaya. 1975a. Accumulation of radionuclides in coastal sediment of Japan (II) Contents of fission products in some coastal sediments collected in 1966-1972. *J. Radiat. Res.* 16:184-192.
- Nakamura, K., and Y. Nagaya. 1975b. Dispersion and accumulation of radionuclides in sediments of Urazoko Bay (I). *J. Oceanogr. Soc. Jpn.* 31:145-153.
- Nakamura, K., and Y. Nagaya. 1985. Accumulation of cesium-137 and plutonium-239, 240 in sediments of the coastal sea and the north Pacific. p. 171-180. *In* A.C. Sigleo and A. Hattori (ed.) *Mar. Estuarine Geochem.* Lewis Publ., Inc., Chelsea, MI.
- †Nelson, J.L., R.W. Perkins, J.M. Neilsen, and W.L. Hausfeld. 1966. Reaction of radionuclides from Hanford reactors with Columbia River sediments. p. 139-161. *In* Disposal of radioactive waste in seas, oceans, and surface waters. STI/PUB/126-139-161. Int. Atomic Energy Agency, Vienna.
- Nishita, H., A.J. Steen, and K.H. Larson. 1958. Release of Sr^{90} and Cs^{137} from Vina loam upon prolonged cropping. *Soil Sci.* 86:195-201.
- Norton, S.A. 1986. A review of the chemical record in lake sediment of energy related air pollution and its effects on lakes. *Water Air*

- Soil Pollut. 30:331-345.
- Nriagu, J.O. 1984. Role of inland water sediments as sinks for anthropogenic sulfur. *Sci. Total Environ.* 38:7-13.
- Nriagu, J.O., A.L.W. Kemp, H.K.T. Wong, and N. Harper. 1979. Sedimentary record of heavy metal pollution in Lake Erie. *Geochim. Cosmochim. Acta* 43:247-258.
- Nyffeler, U.P., Y.-H. Li, and P.H. Santschi. 1984. A kinetic approach to describe trace element distribution between particles and solution in natural aquatic systems. *Geochim. Cosmochim. Acta* 48:1513-1522.
- Officer, C.B. 1982. Mixing sedimentation rates and age dating for sediment cores. *Mar. Geol.* 46:261-278.
- Officer, C.B., and D.R. Lynch. 1982. Interpretation procedures for the determination of sediment parameters from time-dependent flux inputs. *Earth Planet Sci. Lett.* 61:55-62.
- Okuda, S. 1982. Sedimentary environments in Lake Biwa. Estimate of recent sedimentation rate. (In Japanese.) *Kankyo Gijutsu* 11:401-403.
- Oldfield, F. 1977. Lakes and their drainage basins as units of sediment-based ecological study. *Progr. Phys. Geogr.* 1:460-504.
- Oldfield, F., and P.G. Appleby. 1984. A combined radiometric and mineral magnetic approach to recent geochronology in lakes affected by catchment disturbance and sediment redistribution. *Chem. Geol.* 44:67-83.
- Oldfield, F., P.G. Appleby, and R.W. Battarbee. 1978. Alternative ^{210}Pb dating: Results from New Guinea Highlands and Lough Erne. *Nature (London)* 271:339-342.
- Oldfield, F., P.G. Appleby, R.S. Cambray, J.D. Eakins, K.E. Barber, R.W. Battarbee, G.R. Pearson, and J.M. Williams. 1979. ^{210}Pb , ^{137}Cs and ^{239}Pu profiles in ombrotrophic peat. *Oikos* 33:40-45.
- Oldfield, F., P.G. Appleby, and R. Thompson. 1980. Palaeoecological studies of lakes in the highlands of Papua New Guinea. I. The chronology of sedimentation. *J. Ecol.* 68:457-477.
- Oldfield, F., R.W. Battarbee, and J.A. Dearing. 1983. New approaches to recent environmental change. *Geogr. J.* 149:167-181.
- Oldfield, F., and B. Maher. 1984. A mineral magnetic approach to erosion studies. p. 161-168. *In* R.J. Loughran (ed.) *Drainage basin erosion and sedimentation*. Univ. of Newcastle, and Soil Conserv. Serv., N.S.W., Australia.
- Oldfield, F., A.T. Worsley, and P.G. Appleby. 1985. Evidence from lake sediments for recent erosion rates in the highlands of Papua New Guinea. p. 185-196. *In* I. Douglas and T. Spencer (ed.) *Environmental change and tropical geomorphology*. Allen and Unwin, London.
- Olsen, C.R. 1979. Radionuclides, sedimentation and the accumulation of pollutants in the Hudson estuary. Ph.D. thesis. Columbia Univ., New York.
- Olsen, C.R., N.H. Cutshall, and I.L. Larsen. 1982. Pollutant-particle associations and dynamics in coastal marine environments: A review. *Mar. Chem.* 11:501-533.
- Olsen, C.R., N.H. Cutshall, I.L. Larsen, H.J. Simpson, R.M. Trier, and R.F. Bopp. 1985. An estuarine fine-particle budget determined from radionuclide tracers. *Geo-Marine Lett.* 4:157-160.
- Olsen, C.R., I.L. Larsen, N.H. Cutshall, J.F. Donoghue, O.P. Bricker, and H.J. Simpson. 1981a. Reactor-released radionuclides in Susquehanna River sediments. *Nature (London)* 294:242-245.
- Olsen, C.R., H.J. Simpson, T.-H. Peng, R.F. Bopp, and R.M. Trier. 1981b. Sediment mixing and accumulation rates effects on radionuclide depth profiles in Hudson estuary sediments. *J. Geophys. Res.* 86:11020-11028.
- Olsen, C.R., H.J. Simpson, and R.M. Trier. 1981c. Plutonium, radiocesium and radiocobalt in sediments of the Hudson River estuary. *Earth Planet Sci. Lett.* 55:377-392.
- Onenema, O., and R.D. DeLaune. 1988. Accretion rates in salt marshes in the Eastern Scheldt, South-west Netherlands. *Estuarine, Coastal Shelf Sci.* 26:379-394.
- Osborne, P.L., and B. Moss. 1977. Paleolimnology and trends in the phosphorus and iron budgets of an old man-made lake, Barton Broad, Norfolk. *Freshwater Biol.* 7:213-233.
- Parise, G., G. Paleologo, G. Premazzi, P. Pulici, A. Riva, T. Ruaro, R. Bugini, L. De Capitani, and A. Maccagni. 1981. Lake Como (Northern Italy): Mineralogical, chemical and biological characteristics of the sediments. *European Appl. Res. Rep.-Environ. Nat. Res. Sect.* 1(3):385-416.
- Parise, G., and G. Premazzi. 1986. I fenomeni di compattazione nella geocronologia dei sedimenti lacustri. (In Italian.) *Acqua-Aria* 8:783-790.
- Parker, J.I., and D.N. Edgington. 1976. Concentration of diatom frustules in Lake Michigan sediment cores. *Limnol. Oceanogr.* 21:887-893.
- Patel, B., S. Patel, and S. Pawar. 1978. Desorption of radioactivity from nearshore sediments. *Estuarine Coastal Mar. Sci.* 7:49-58.
- Peart, M.R., and D.E. Walling. 1986. Fingerprinting sediment source: The example of a drainage basin in Devon, U.K. p. 41-55. *In* Drainage basin sediment delivery. Proc. of the Albuquerque Symposium, Albuquerque, NM. 4-8 August. IAHS Publ. 159. IAHS Press, Wallingford, U.K.
- Peart, M.R., and D.E. Walling. 1988. Techniques for establishing suspended sediment sources in two drainage basins in Devon, U.K.: A comparative assessment. IAHS Publ. 174:269-279.
- Pegoyev, A.N., and Sh.D. Fridman. 1978. Vertical profiles of cesium-137 in soils. (Translated.) *Pochvovedeniye* 8:77-81.
- Peirson, D.H., R.S. Cambray, P.A. Cawse, J.D. Eakins, and N.J. Pattenden. 1982. Environmental radioactivity in Cumbria. *Nature (London)* 300:27-31.
- Pennington, W. 1973. The recent sediments of Windermere. *Freshwater Biol.* 3:363-382.
- Pennington, W. 1974. Seston and sediment formation in five Lake District lakes. *J. Ecol.* 62:215-251.
- Pennington, W. 1978. Responses of some British lakes to past changes in land use on their catchments. *Verh. Int. Verein. Limnol.* 20:636-641.
- Pennington, W. 1979. The origin of pollen in lake sediments: An enclosed lake compared with one receiving inflow streams. *New Phytol.* 83:189-213.
- Pennington, W. 1981. Records of a lake's life in time: The sediments. *Hydrobiologia* 79:197-219.
- Pennington, W., R.S. Cambray, J.D. Eakins, and D.D. Harkness. 1976. Radionuclide dating of the recent sediments of Blelham Tarn. *Freshwater Biol.* 6:317-331.
- Pennington, W., R.S. Cambray, and E.M. Fisher. 1973. Observations on lake sediments using fallout ^{137}Cs as a tracer. *Nature (London)* 242:324-326.
- Pennock, D.J., and E. DeJong. 1987. The influence of slope curvature on soil erosion and deposition in hummock terrain. *Soil Sci.* 144:209-217.
- Perkins, R.W., and C.W. Thomas. 1980. Worldwide fallout. p. 53-82. *In* W.C. Hanson (ed.) *Transuranic elements in the environment*. USDOE/TIC-22800. US DOE, Washington, DC.
- Peterson, S.A. 1982. Lake restoration by sediment removal. *Water Resour. Bull.* 18:423-435.
- Petit, D., M. Thomas, and L. Lamberts. 1987. Origin of heavy metal fluxes to the Meuse River in Southern Belgium using ^{210}Pb -dated water-meadow sediments. *J. Environ. Radioactivity* 5:303-316.
- Phillips, F.M., G.I. Smith, H.W. Bentley, D. Elmore, and H.E. Gove. 1983. Chlorine-36 dating of saline sediments: Preliminary results from Searles Lake, California. *Science (Washington, DC)* 222:925-927.
- Pickering, R.J. 1969. Distribution of radionuclides in bottom sediment at the Clinch River, eastern Tennessee. U.S. Geol. Surv. Prof. Pap. 433-H.
- Pickering, R.J., P.H. Carrigan, Jr., and F.L. Parker. 1965. The Clinch River Study—an investigation of the fate of radionuclides released to a surface stream. U.S. Geol. Surv. Circ. 497.
- Pickering, R.J., P.H. Carrigan, Jr., T. Tamura, H.H. Abee, J.W. Beverage, and R.W. Andrews, Jr. 1966. Radioactivity in the bottom sediment of the Clinch and Tennessee Rivers. p. 57-86. *In* Disposal of radioactive wastes into seas, oceans and surface waters: Vienna, Austria. Int. Atomic Energy Agency, Vienna.
- Pinder, J.E., III, C.T. Garten, Jr., and D. Paine. 1980. Factors affecting radiocesium uptake by plants inhabiting a contaminated floodplain. *Acta Ecologica* 1:3-10.
- Plato, P. 1974. Use of rivers to predict accumulation in sediments of radionuclides discharged from nuclear power stations. *Health Phys.* 26:489-496.
- Plato, P. 1975. Letter to editor. *Health Phys.* 28:635-636.
- Plato, P., and G.C. Goldman. 1972. Use of fallout cesium-137 as a tracer to define the recent deltaic facies of a river. *Radiat. Data Rep.* 13:653-657.
- Plato, P.A. 1972. Distribution of cesium-137 and naturally occurring radionuclides in sediments of Lake Michigan. *Radiat. Data Rep.* 13:181-187.
- Plato, P.A., and A.P. Jacobson. 1976. Cesium-137 in Lake Michigan sediment: Areal distribution and correlation with other man-made materials. *Environ. Pollut.* 10:19-34.
- Popp, C.J., J.W. Hawley, D.W. Love, and M. Dehn. 1988. Use of radiometric (Cs-137 , Pb-210), geomorphic, and stratigraphic techniques to date recent oxbow sediments in the Rio Puerco drainage Grants uranium region, New Mexico. *Environ. Geol. Water Sci.* 11:253-269.
- Powers, C.F., F.S. Stay, and W.D. Sanville. 1975. Aquatic sediments. *J. Water Pollut. Control Fed.* 47:1611-1617.
- Premazzi, G. 1978. Metodo di valutazione della rate di sedimentazione. *Atti del Convegno sulla "Eutrofizzazione in Italia"*, Roma, 3-4 Ottobre 1978. AC/2/55. p. 181-195. (In Italian.)
- Premazzi, G. 1979. The Cs^{137} technique to evaluate recent sedimentation rates. p. 181-196. *In* R. Marchetti (ed.) *Convegno Sulla Eutrofizzazione in Italia*. AC/2/45-70, Roma, Italy.

- Premazzi, G., and G. Marengo. 1982. Sedimentation rates in a Swiss-Italian lake measured with sediment traps. *Hydrobiologia* 92:603-610.
- Premazzi, G., A. Provini, G.F. Gaggino, and G. Parise. 1984. Geochemical trends in sediments from 13 Italian subalpine lakes. p. 157-165. In P.G. Sly (ed.) *Sediment and water interactions*. Springer-Verlag, New York.
- Premazzi, G., and O. Ravera. 1977. Chemical characteristics of Lake Lugano sediments. p. 121-124. In H.L. Golterman (ed.) *Interactions between sediments and fresh water*. Dr. W. Junk B.V. Publ., The Hague, Netherlands.
- Prokhorov, V.M. 1975. Forecasting Cs^{137} migration in soils. (Translated.) *Pochvovedeniye* 11:60-67.
- Provini, A., and G.F. Gaggino. 1984a. Depth profiles of Cu, Cr, and Zn in Lake Orta sediments (Northern Italy). p. 167-174. In P.G. Sly (ed.) *Sediment and water interactions*. Springer-Verlag, New York.
- Provini, A., and G.F. Gaggino. 1984b. Metal contamination and historical records in Lake Orta sediments (Northern Italy). p. 318-321. In Proc. 3rd Int. Symp. on the Interactions between Sediments and Water, Geneva.
- Provini, A., and G.F. Gaggino. 1985. Sediment as a record of copper pollution in Lake Orta. *Verh. Int. Verein. Limnol.* 22:2390-2393.
- Provini, A., G. Premazzi, and G.F. Gaggino. 1987a. Valutazione del rischio potenziale da metalli pesanti nei laghi mediante l'analisi dei sedimenti. (In Italian.) *Ing. Ambientale* 16:68-76.
- Provini, A., G. Premazzi, S. Galassi, and G.F. Gaggino. 1987b. Distribution of nutrients, trace elements PAHs, and radionuclides in sediment cores from Lake of Varese (N. Italy). Proc. of 4th Int. Symp. in Interaction between Sediments and Water, Melbourne, Australia.
- Rapin, F. 1981. Chronologie et evolution de la contamination par les metaux lourds des sediments de la Baie des Anges (Mediterranee, France). *Environ. Technol. Lett.* 2:253-262. (French).
- †Ravera, O. 1961. Sediments. p. 31-37. In *Accumulation of fission products from fall-out in Lake Biota (Lake Maggiore)*. Final Rep. CNI-109. Int. Atomic Energy Agency, Vienna. IAEA Contract 59.
- †Ravera, O., and G. Premazzi. 1971. A method to study the history of any persistent pollution in a lake by the concentration of ^{137}Cs from fallout. *Publ. Biol. Directorate, Comm. European Communities* 679:1-16.
- Ravera, O., and G. Premazzi. 1972. A method to study the history of any persistent pollution in a lake by the concentration of ^{137}Cs from fallout in sediments. p. 703-722. In *Symp. on radioecology applied to the protection of man and his environment*. Vol. I. EUR 4800. Int. Atomic Energy Agency, Vienna.
- Ravera, O., and M. Viola. 1977. Sedimentation-rate in a basin (Agno) of Lake Lugano. p. 174-178. In H.L. Golterman (ed.) *Interactions between sediment and fresh water*. Dr. W. Junk B.V. Publ., The Hague, Netherlands.
- Ravera, O., and S. Zarini. 1987. A century of variations in some zooplankton and zoobenthos organisms in Lake Comabbio (Northern Italy) 'read' from their remains in the lake sediment. *Schweiz. Z. Hydrol.* 49:93-101.
- Rea, D.K., R.A. Bourbonniere, and P.A. Meyers. 1980. Southern Lake Michigan sediments: Changes in accumulation rate, mineralogy, and organic content. *J. Great Lakes Res.* 6:321-330.
- Ridgeway, I.M., and N.B. Price. 1987. Geochemical associations and post-depositional mobility of heavy metals in coastal sediments: Loch Etive, Scotland. *Mar. Chem.* 21:229-248.
- Riel, G.K. 1970. Radioactive cesium in estuaries. *Radiol. Health Data Rep.* 11:659-665.
- Ritchie, J.C. 1988. Organic matter content in sediments of three navigation pools along the upper Mississippi River. *J. Freshwater Ecol.* 4:343-349.
- Ritchie, J.C. 1989. Carbon content of sediments in agricultural reservoirs. *Water Resour. Bull.* 25:301-308.
- †Ritchie, J.C., E.E.C. Clebsch, and W.K. Rudolph. 1970a. Distribution of fallout and natural gamma radionuclides in litter, humus, and surface mineral soils under natural vegetation in the Great Smoky Mountains, North Carolina-Tennessee. *Health Phys.* 18:479-491.
- Ritchie, J.C., C.M. Cooper, and J.R. McHenry. 1979. Recent accumulation of sediment in lakes in the Bear Creek watershed in the Mississippi Delta. *Southeast. Geol.* 20:173-180.
- Ritchie, J.C., C.M. Cooper, and J.R. McHenry. 1986. Sediment accumulation rates in lakes and reservoirs in the Mississippi River Valley. p. 1357-1365. In S.Y. Wang et al. (ed.) *River sedimentation*. Vol. III. The Univ. of Mississippi, Oxford, MS.
- Ritchie, J.C., C.M. Cooper, J.R. McHenry, and F.R. Schiebe. 1983. Sediment accumulation in Lake Chicot, Arkansas. *Environ. Geol.* 5:79-82.
- †Ritchie, J.C., P.H. Hawks, and J.R. McHenry. 1975. Deposition rates in valleys determined using fallout cesium-137. *Geol. Soc. Am. Bull.* 86:1128-1130.
- †Ritchie, J.C., and J.R. McHenry. 1973a. Vertical distribution of fallout cesium-137 in cultivated soils. *Radiat. Data Rep.* 14:727-728.
- †Ritchie, J.C., and J.R. McHenry. 1973b. Determination of fallout ^{137}Cs and naturally occurring gamma-ray emitters in sediments. *Intl. J. Appl. Radiat. Isotopes* 24:575-578.
- †Ritchie, J.C., and J.R. McHenry. 1975a. Fallout Cs-137: A tool in conservation research. *J. Soil Water Conserv.* 30:283-286.
- Ritchie, J.C., and J.R. McHenry. 1975b. Fallout cesium-137 in estuarine sediments. *J. Miss. Acad. Sci.* 20:34-39.
- †Ritchie, J.C., and J.R. McHenry. 1976. Fallout ^{137}Cs in soils and sediments of some Texas watershed ecosystems. p. 229-303. In C.C. Cushing, Jr. (ed.) *Radioecology and energy resources*. Ecol. Soc. Am. Spec. Publ. I. Dowden, Hutchinson, and Ross, Inc., Stroudsburg, PA.
- Ritchie, J.C., and J.R. McHenry. 1977a. The distribution of Cs-137 in some watersheds in the eastern United States. *Health Phys.* 32:101-105.
- Ritchie, J.C., and J.R. McHenry. 1977b. A rapid method for determining recent deposition rates of freshwater sediments. p. 203-207. In H.L. Golterman (ed.) *Interaction between sediments and fresh water*. Dr. W. Junk B.V. Publ., The Hague, Netherlands.
- Ritchie, J.C., and J.R. McHenry. 1978. Fallout cesium-137 in cultivated and noncultivated north central United States watersheds. *J. Environ. Qual.* 7:40-44.
- Ritchie, J.C., and J.R. McHenry. 1982. Redistribution of fall-out caesium-137 in small watersheds in the United States of America. p. 809-810. In *Environmental migration of long-lived radionuclides*. IAEA-SM-257/68P. Int. Atomic Energy Agency, Vienna.
- Ritchie, J.C., and J.R. McHenry. 1984. Cesium-137 and sediment deposition. p. 183-188. In R.J. Loughran (ed.) *Drainage basin erosion and sedimentation*. Univ. of Newcastle, and Soil Conserv. Serv. of N.S.W., Australia.
- Ritchie, J.C., and J.R. McHenry. 1985. A comparison of three methods of measuring recent rates of sediment accumulation. *Water Resour. Res.* 21:99-103.
- †Ritchie, J.C., J.R. McHenry, and G.D. Bubenzer. 1982. Redistribution of fallout ^{137}Cs in Brunner Creek watershed in Wisconsin. *Wis. Acad. Sci. Arts Lett.* 70:161-166.
- Ritchie, J.C., J.R. McHenry, and A.C. Gill. 1970b. Distribution of fallout cesium-137 in the upper 4-inches of soil in relation to vegetation type. *ASB Bull.* 17(2):60.
- Ritchie, J.C., J.R. McHenry, and A.C. Gill. 1972a. Dating recent reservoir sediment using radioactive fallout. *ASB Bull.* 19(2):95.
- †Ritchie, J.C., J.R. McHenry, and A.C. Gill. 1972b. The distribution of ^{137}Cs in the upper 10 cm of soil under different cover types in northern Mississippi. *Health Phys.* 22:197-198.
- †Ritchie, J.C., J.R. McHenry, and A.C. Gill. 1973. Dating recent reservoir sediments. *Limnol. Oceanogr.* 18:254-263.
- †Ritchie, J.C., J.R. McHenry, and A.C. Gill. 1974. Fallout ^{137}Cs in the soils and sediments of three small watersheds. *Ecology* 55:887-890.
- †Ritchie, J.C., J.R. McHenry, A.C. Gill, and P.H. Hawks. 1970c. The use of fallout cesium-137 as a tracer of sediment movement and deposition. p. 149-162. In *Proc. Miss. Water Resour. Conf.*, Jackson, MS. 15 April. Mississippi Water Resour. Inst., Mississippi State, MS.
- †Ritchie, J.C., J.R. McHenry, A.C. Gill, and P.H. Hawks. 1970d. Distribution of fallout cesium-137 in sediment profiles. *Health Phys.* 19:334.
- †Ritchie, J.C., J.R. McHenry, A.C. Gill, and P.H. Hawks. 1971. Distribution of cesium-137 in a small watershed in northern Mississippi. p. 129-132. In *Proc. Third national symposium on radioecology conf-710-501-Pl*, Oak Ridge, TN. 10-12 May. U.S. Atomic Energy Comm., Washington, DC.
- Ritchie, J.C., J.R. McHenry, A.C. Gill, and P.H. Hawks. 1972. Fallout ^{137}Cs in reservoir sediment. *Health Phys.* 22:97-98.
- †Ritchie, J.C., J.A. Spraberry, and J.R. McHenry. 1974. Estimating soil erosion from the redistribution of fallout ^{137}Cs . *Soil Sci. Soc. Am. Proc.* 38:137-139.
- Robbe, D., P. Marchandise, and A. Thomas. 1983. Determination of the origin of metallic pollutants by dating of sediment cores. (In French.) *Bull. Liaison Lab. Ponts. Chaussees* 127:81-92.
- Robbins, J.A. 1978. Geochemical and geophysical applications of radioactive lead. p. 285-393. In J.O. Nriagu (ed.) *The biochemistry of lead in the environment*. Elsevier/North Holland Biomedical Press, Amsterdam.
- Robbins, J.A. 1980. Sediments of southern Lake Huron: Elemental composition and accumulation rates. *USEPA, Ecol. Res. Ser. EPA-600/3-80-080*. USEPA, Washington, DC.
- Robbins, J.A. 1982. Stratigraphic and dynamic effects of sediment reworking by Great Lakes zoobenthos. *Hydrobiologia* 92:611-622.

- Robbins, J.A. 1985a. Great Lakes regional fallout source functions. NOAA Tech. Memo. ERL GLERL-56. NOAA, Ann Arbor, MI.
- Robbins, J.A. 1985b. The Coupled Lake Model for estimating the long-term response of the Great Lakes to time-dependent loadings of particle-associated contaminants. NOAA Tech. Memo. ERL GLERL-57. NOAA, Ann Arbor, MI.
- †Robbins, J.A. 1986a. A model for particle-selective transport of tracers in sediments with conveyor belt deposit feeders. *J. Geophys. Res.* 91:8542-8558.
- Robbins, J.A. 1986b. Sediments of Saginaw Bay, Lake Huron: Elemental composition and accumulation rates. Great Lakes and Marine Waters Center, Spec. Rep. 102 of the Great Lakes Res. Div., Univ. of Michigan, Ann Arbor, MI.
- Robbins, J.A., E. Callender, and Nancy R. Morehead. 1988. Records of fallout cesium-137 in Lake Oahe Reservoir, South Dakota. *EOS* 69:68.
- †Robbins, J.A., and D.N. Edgington. 1972. The use of natural and fallout radionuclides to measure recent sedimentation rates in Southern Lake Michigan. Argonne Natl. Lab. Rep. ANL-7960. Part 3. p. 31-53. Argonne Natl. Lab., Argonne, IL.
- Robbins, J.A., and D.N. Edgington. 1974. The distribution of trace elements in the clay silt sediments of Grand Haven. p. 78-85. *In* Radiological and Environ. Res. Div. Annu. Rep., Ecology. Argonne Natl. Lab., Argonne, IL.
- †Robbins, J.A., and D.N. Edgington. 1975. Determination of recent sedimentation rates in Lake Michigan using Pb-210 and Cs-137. *Geochim. Cosmochim. Acta* 39:285-304.
- Robbins, J.A., and D.N. Edgington. 1976. Depositional processes and the determination of recent sediment rates in Lake Michigan. p. 378-390. *In* Proc. Second Federal Conf. on the Great Lakes, Argonne, IL. 25-27 March. Great Lakes Basin Comm., Ann Arbor, MI.
- Robbins, J.A., D.N. Edgington, and A.L.W. Kemp. 1977a. Comparative lead-210, cesium-137 and pollen geochronologies of recent sediments from Lakes Erie and Ontario. p. 87-92. *In* Radiological and Environ. Res. Div. Annu. Rep., Ecology. Argonne Natl. Lab., Argonne, IL.
- †Robbins, J.A., D.N. Edgington, and A.L.W. Kemp. 1978. Comparative ^{210}Pb , ^{137}Cs , and pollen geochronologies of sediments from Lakes Ontario and Erie. *Quat. Res. (NY)* 10:256-278.
- Robbins, J.A., K. Husby-Coupland, and D.S. White. 1984. Precise radiotracer measurement of the rate of sediment reworking by *Stylodrilus heringianus* and the effects of variable dissolved oxygen concentrations. *J. Great Lakes Res.* 10:335-347.
- Robbins, J.A., T. Keilty, D.S. White, and D.N. Edgington. 1989. Relationship among Tubificid abundances, sediment composition, and accumulation rates in Lake Erie. *Can. J. Fish. Aquat. Sci.* 46:223-231.
- †Robbins, J.A., J.R. Krezoski, and S.C. Mozley. 1977b. Radioactivity in sediments of the Great Lakes: Post-depositional redistribution by deposit-feeding organisms. *Earth Planet. Sci. Lett.* 36:325-333.
- †Robbins, J.A., P.L. McCall, J.B. Fisher, and J.R. Krezoski. 1979. Effect of deposit feeders on migration of ^{137}Cs in lake sediments. *Earth Planet. Sci. Lett.* 42:277-287.
- Robbins, J.A., and C.L. Schelske. 1988. Biogenic silica deposition in Lake Erie since 1800. *EOS Trans.* 69:1143.
- †Rogowski, A.S., and T. Tamura. 1965. Movement of ^{137}Cs by runoff, erosion and infiltration on the alluvial Captina silt loam. *Health Phys.* 11:1333-1340.
- †Rogowski, A.S., and T. Tamura. 1970a. Environmental mobility of cesium-137. *Radiat. Bot.* 10:35-45.
- †Rogowski, A.S., and T. Tamura. 1970b. Erosional behavior of cesium-137. *Health Phys.* 18:467-477.
- Rose, C.W., M.E. McCallan, B.M. O'Leary, and G. Sander. 1980. Testing a model of field-scale soil erosion and deposition on a plane land element using the fallout isotope ^{137}Cs . The Inst. of Eng., Australia, Conf. on Agric. Eng., Geelong, Reprint of Papers 80/7. Inst. of Eng., Australia.
- Rummery, T.A., J. Bloemendal, J. Dearing, F. Oldfield, and R. Thompson. 1979. The persistence of fire-induced magnetic oxides in soils and lake sediments. *Ann. Geophys.* 35:103-107.
- Salo, A., R. Saxen, and M. Puhakainen. 1984. Transport of airborne ^{90}Sr and ^{137}Cs deposited in the basins of the five largest rivers in Finland. *Aqua Fenn.* 14:21-31.
- Salo, A., K. Tuomainen, and A. Voipio. 1986. Inventories of some long-lived radionuclides in the Baltic Sea. *Sci. Total Environ.* 54:247-260.
- Santschi, P. 1984. Radionuclides as tracers for sedimentation and remobilization processes in the ocean and in lakes. p. 437-449. *In* P.G. Sly (ed.) Sediments and water interactions. Springer-Verlag, New York.
- Santschi, P.H. 1987. Chernobyl radionuclides in the environment: Tracers for the tight coupling between atmospheric, terrestrial and aquatic geochemical processes. *EAWAG News* 22/23:1-6.
- Santschi, P.H., S. Bollhalder, K. Farrenkoth, A. Lueck, S. Zingg, and M. Sturm. 1988. Chernobyl radionuclides in the environment: Tracers for the tight coupling between atmospheric, terrestrial and aquatic geochemical processes. *Environ. Sci. Technol.* 22:510-516.
- Santschi, P.H., Y.-H. Li, D.M. Adler, M. Amdurer, J. Bell, and U.P. Nyffeler. 1983. The relative mobility of natural (Th, Pb, Po) and fallout (Pu, Am, Cs) radionuclides in the coastal marine environments: Results from model ecosystems (MERL) and Narragansett Bay. *Geochim. Cosmochim. Acta* 47:201-210.
- Santschi, P.H., Y.-H. Li, and S. Carson. 1980. The fate of trace metals in Narragansett Bay, Rhode Island: Radiotracer experiments in microcosms. *Estuarine Coastal Mar. Sci.* 10:635-654.
- Santschi, P.H., Y.-H. Li, P. O'Hara, M. Amdurer, D. Adler, and P. Doering. 1987. The relative mobility of radioactive trace elements across the sediment-water interface of the MERL model ecosystems of Narragansett Bay. *J. Mar. Res.* 45:1007-1048.
- Santschi, P.H., S. Nixon, M. Pilson, and C. Hunt. 1984. Accumulation of sediments, trace metals (Pb, Cu) and hydrocarbons in Narragansett Bay, Rhode Island. *Estuarine Coastal Shelf Sci.* 19:427-450.
- Santschi, P.H., U.P. Nyffeler, R.F. Anderson, S.L. Schiff, and P. O'Hara. 1986. Response of radioactive trace metals to acid-base titrations in controlled experimental ecosystems: Evaluation of transport parameters for application to whole lake radiotracer experiments. *Can. J. Fish. Aquat. Sci.* 43:60-77.
- Santschi, P.H., and P.W. Schindler. 1977. Chemical and geochemical studies of Lake Biel, I. A mass balance for Lake Biel and its implications for the rates of erosion in the drainage area. *Swiss. J. Hydrol.* 39:181-200.
- Santschi, P.H., C. Schuler, M. Strum, A. Lueck, K. Farrenkoth, S. Bollhalder, and E. Wieland. 1988. Natural radionuclides in Lake Zurich: Tracers for particle and tracer element dynamics. *EOS Trans.* 69:1085.
- Schimmack, W., K. Bunzl, and H. Bachhuber. 1987. Variability of the sorption of Cs, Zn, Sr, Co, Cd, Ce, Ru, Tc, and I at trace concentrations by a forest soil along a transect. *Environ. Int.* 13:427-436.
- Schoenburg, M. 1987. Radiometric dating and quantitative analysis of elements in depth profiles of sediments by means of nuclear physical as well as x-ray fluorescence and atomic emission spectroscopic methods. Fachbereich 12-Physik., Rep. GKSS-87/E/54. Hamburg Univ. (In German.)
- Schreiber, B. 1967a. Ecology of Acantharia in relation to Sr circulation in the sea. Final Rep. IAEA Contract US/62. Int. Atomic Energy Agency, Vienna.
- Schreiber, B. 1967b. Radionuclides in marine plankton and in coastal sediments. p. 753-770. *In* B. Aberg and F.P. Hungate (ed.) Radioecological concentration processes. Pergamon Press, New York.
- †Schreiber, B. 1968. Essay of a method of absolute dating of the coastal marine sediments by means of the vertical distribution of the fallout radionuclides. *Accademia Nazionale Dei Lincei, Serie VII, Vol. XLV, Fasc. 6*:251-257.
- Schreiber, B. 1969. Significance of sediments in evaluating the radioactive contamination of the sea. Fourth Int. Colloq. Medical Oceanography, Naples, Italy.
- Schreiber, B. 1974. Plankton and sediment: First and last step of the radioactivity diffusion in the area. *Ciencia Cult. (San Paulo)* 27:197-206.
- Schreiber, B., E. Cerrai, C. Triulzi, L.T. Pelati, and M. Mezzadri. 1969. Ricerche sulla distribuzione verticale della radioattività nei sedimenti del Mar Ligure. (In Italian.) *G. Fis. Sanit. Prot. Radiaz.* 13:1-19.
- Schreiber, B., L.T. Pelati, E. Cerrai, and C. Triulzi. 1964. Gross beta radioactivity of littoral sediments of the Ligurian Sea. *Energ. Nucl. (Milan)* 11:616-624.
- †Schreiber, B., L.T. Pelati, M.G. Mezzadri, and G. Motta. 1968. Gross beta radioactivity in sediments of the North Adriatic Sea: A possibility of evaluating the sedimentation rate. *Arch. Oceanogr. Limnol.* 16:45-62.
- Schroeder, R.A. 1985. Sediment accumulation rates in Irondequoit Bay, New York based on lead-210 and cesium-137 geochronology. *Northeast. Environ. Sci.* 4:23-29.
- †Schulz, R.K. 1965. Soil chemistry of radionuclides. *Health Phys.* 11:1317-1324.
- †Schulz, R.K., R. Overstreet, and I. Barshad. 1960. On the soil chemistry of cesium 137. *Soil Sci.* 89:19-27.
- †Sharma, P., L.R. Gardner, W.S. Moore, and M.S. Bollinger. 1987. Sedimentation and bioturbation in a salt marsh as revealed by ^{210}Pb , ^{137}Cs , and ^7Be studies. *Limnol. Oceanogr.* 32:313-326.
- Sholkovitz, E.R. 1985. Redox-related geochemistry in lakes: Alkali metals, alkaline-earth elements, and ^{137}Cs . p. 119-142. *In* W.

- Stumm (ed.) Chemical processes in lakes. John Wiley & Sons, New York.
- Sholkovitz, E.R., J.K. Cochran, and A.E. Carey. 1983. Laboratory studies of the diagenesis and mobility of $^{239,240}\text{Pu}$ and ^{137}Cs in nearshore sediments. *Geochim. Cosmochim. Acta* 47:1369-1379.
- Sholkovitz, E.R., and D.R. Mann. 1984. The pore water chemistry of $^{239,240}\text{Pu}$ and ^{137}Cs in sediments of Buzzards Bay, Massachusetts. *Geochim. Cosmochim. Acta* 48:1107-1114.
- Silantyev, A.N., I.G. Shkuratova, and R.N. Khatskevich. 1978. Spatial distribution of cesium-137 in the soils of the European USSR. *Soviet Soil Sci.* 10(2):187-188. (English Translation.) *Pochvovedeniye* (1978) 4:47-48.
- Simpson, H.J., C.R. Olsen, R. Bopp, R.M. Trier, and S.C. Williams. 1978. Cesium-137 as a tracer for reactive pollutants in estuarine sediments. First American-Soviet Symposium on Chemical Pollution of Marine Environ. USEPA Rep. 600/9-78-038. USEPA, Gulf Breeze, FL.
- Simpson, H.J., C.R. Olsen, R.M. Trier, and S.C. Williams. 1976. Man-made radionuclides and sedimentation in the Hudson River estuary. *Science* (Washington, DC) 194:179-183.
- Simpson, H.J., S.C. Williams, C.R. Olsen, and D.E. Hammond. 1977. Nutrient and particulate matter budgets in urban estuaries. p. 94-103. *In* Estuaries, geophysics and the environment. Natl. Acad. of Sci., Washington, DC.
- Skagius, K., and I. Neretnieks. 1988. Measurements of cesium and strontium diffusion in biotite gneiss. *Water Resour. Res.* 24:75-84.
- Smith, C.J., R.D. DeLaune, and W.H. Patrick, Jr. 1983. Carbon dioxide emission and carbon accumulation in coastal wetlands. *Estuarine Coastal Shelf Sci.* 17:21-29.
- Smith, J.N., and K.M. Ellis. 1982. Transport mechanism for Pb-210, Cs-137 and Pu fallout radionuclides through fluvial-marine systems. *Geochim. Cosmochim. Acta* 46:941-954.
- Smith, J.N., K.M. Ellis, and D.M. Nelson. 1987. Time-dependent modeling of fallout radionuclide transport in a drainage basin: Significance of "slow" erosional and "fast" hydrological components. *Chem. Geol.* 63:157-180.
- Smith, J.N., and C.T. Schafer. 1987. A 20th-century record of climatologically modulated sediment accumulation rates in a Canadian fjord. *Quat. Res. (NY)* 27:232-247.
- Smith, J.N., and A. Walton. 1980. Sediment accumulation rates and geochronologies measured in the Saguenay Fjord using Pb-210 dating method. *Geochim. Cosmochim. Acta* 44:225-240.
- Smith, S.J., J.R. McHenry, R.G. Menzel, and N.H. Welch. 1986. Agricultural chemicals and heavy metals in upland soils and valley alluviums in the Little Washita River basin. *J. Soil Water Conserv.* 41:333-336.
- Soszka, G.J., D. Grzybowska, J. Rostek, A. Pietruszewski, T. Wardaszko, A. Kalinowska, and J. Tomczak. 1986. Radioecological characteristics of Lake Zarnowieckie. *Pol. Ecol. Studies* 12:363-381.
- †Sprayberry, J.A., and A.J. Bowie. 1969. Predicting sediment yield from complex watersheds. *Trans. Am. Soc. Agric. Eng.* 12:199-201.
- †Spomer, R.G., J.R. McHenry, and R.F. Piest. 1985. Sediment movement and deposition using cesium-137 tracer. *Trans. Am. Soc. Agric. Eng.* 28:767-772.
- Sprugel, D.G., and G.E. Bartelt. 1978. Erosional removal of fallout plutonium from a large midwestern watershed. *J. Environ. Qual.* 7:175-177.
- Stanners, D.A., and S.R. Aston. 1981a. An improved method of determining sedimentation rates by use of artificial radionuclides. *Estuarine Coastal Shelf Sci.* 13:101-106.
- Stanners, D.A., and S.R. Aston. 1981b. Factors controlling the interaction of Cs-137 with suspended and deposited sediments in estuarine and coastal environments. p. 131-141. *In* Impacts of radionuclides releases to the marine environment. IAEA, Vienna.
- Stanners, D.A., and S.R. Aston. 1981c. ^{134}Cs : ^{137}Cs and ^{106}Ru : ^{137}Cs ratios in intertidal sediments from the Cumbria and Lancashire coasts, England. *Estuarine Coastal Shelf Sci.* 13:409-417.
- Stanners, D.A., and S.R. Aston. 1984. The use of reprocessing effluent radionuclides in the geochronology of recent sediments. *Chem. Geol.* 44:19-32.
- Steinberg, C., E. Hammerle, and J. Meihamn. 1983. Changes in the sedimentary fluxes in Lake Walchensee by anthropogenically caused alterations in its catchment area. *Z. Wasser A.B.* 16:48-54.
- Stewart, W.D.P., T. Preston, H.G. Peterson, and N. Christofi. 1982. Nitrogen cycling in eutrophic freshwaters. *Phil. Trans. R. Soc. London B296*:491-509.
- †Stiller, M., and G. Assaf. 1973. Sedimentation and transport of particles in Lake Kinneret traced by ^{137}Cs . *Hydrol. Lakes Symp. IASH Publ. No.* 109:397-403.
- Stiller, M., and D.M. Imboden. 1986. ^{210}Pb in Lake Kinneret waters and sediments: Residence times and fluxes. p. 501-511. *In* P.G. Sly (ed.) Sediment and water interactions. Springer-Verlag, New York.
- Stiller, M., N.E. Yanaki, and J. Kronfeld. 1985. Comparative isotope study of two short sediment cores from the Dead Sea. *Chem. Geol.* 58:107-109.
- Stott, A.P. 1987. Medium-term effects of afforestation on sediment dynamics in a water supply catchment: A mineral magnetic interpretation of reservoir deposits in the Macclesfield Forest, N. W. England. *Earth Surface Processes Landforms* 12:619-630.
- Stow, C.A., R.D. De Laune, and W.H. Patrick, Jr. 1985. Nutrient fluxes in a eutrophic coastal Louisiana freshwater lake. *Environ. Manage.* 9:243-252.
- Svoboda, J., and H.W. Taylor. 1979. Persistence of cesium-137 in Arctic lichens, *Dryas integrifolia*, and lake sediments. *Arct. Alp. Res.* 11:95-108.
- Swan, D.S., M.S. Baxter, I.G. McKinley, and W. Jack. 1982. Radiocaesium and ^{210}Pb in Clyde Sea Loch sediments. *Eстуarine Coastal Shelf Sci.* 15:515-536.
- †Tamura, T. 1964a. Reaction of cesium-137 and strontium-90 with soil minerals and sesquioxides. p. 465-478. *In* 8th Int. Congr. of Soil Sci., Bucharest, Romania. 31 August-9 September. Int. Soc. of Soil Sci., Wageningen, Netherlands.
- †Tamura, T. 1964b. Selective sorption reaction of caesium with mineral soils. *Nucl. Saf.* 5:262-268.
- †Tamura, T., and D.G. Jacobs. 1960. Structural implications in cesium sorption. *Health Phys.* 2:391-398.
- Thomas, M., D. Petit, and L. Lamberts. 1984. Pond sediments as historical record of heavy metal fallout. *Water Air Soil Pollut.* 23:51-59.
- Thorbjarnarson, K.W., C.A. Nittrouer, and D.J. DeMaster. 1986. Accumulation of modern sediment in Quinault Submarine Canyon. *Mar. Geol.* 71:107-124.
- Torgerson, T., A.R. Chivas, and A. Chapman. 1983. Chemical and isotopic characterization and sedimentation rates in Princess Charlotte Bay, Queensland. *BRM J. Aust. Geol. Geophys.* 8:191-200.
- †Torgerson, T., and M.E. Longmore. 1984. ^{137}Cs diffusion in highly organic sediment of Hidden Lake, Fraser Island, Queensland. *Aust. J. Mar. Freshwater Res.* 35:537-548.
- Tracy, B.L., and F.A. Prantl. 1983. 25 years of fission product input to Lakes Superior and Huron. *Water Air Soil Pollut.* 19:15-27.
- Turner, R.R., and S.E. Lindberg. 1978. Behavior and transport of mercury in river-reservoir system downstream of inactive chlor-alkali plant. *Environ. Sci. Technol.* 12:918-923.
- Vakulovskii, S.M., Y.V. Krasnopevtsev, A.I. Nikitin, and V.B. Chumichev. 1982. Distribution of cesium-137 and strontium-90 between the water and bottom sediments in the Black Sea, USSR in 1977. *Okeanologiya* 22:966-969.
- Vakulovskii, S.M., A.I. Nikitin, and V.B. Chumichev. 1983. Cesium-137 and strontium-90 in bottom sediments of the Baltic Sea in 1980. (In Russian.) *Okeanologiya* 23:984-989.
- Van Sickle, J., W.C. Weimer, and D.P. Larsen. 1983. Mixing rates in Shagawa Lake, Minnesota, sediments as determined from ^{106}Ru profiles. *Geochim. Cosmochim. Acta* 47:2189-2197.
- Vaz Carreiro, M.C., and M.M. Sequeria. 1987. Cesium-137 in the Portuguese Rivers Douro and Tejo. *J. Environ. Radioactivity* 5:363-377.
- Vernet, J., E. Davaud, M. Cosandey, and J. Bertie. 1975. Analyse de la radioactivite gamma des sediments des lacs Leman et de Morat. (In French.) *Ecolog. Geol. Helv.* 68:87-96.
- Vernet, J.P., J. Dominik, and P.Y. Favarger. 1984. Texture and sedimentation rates in Lake Geneva. *Environ. Geol.* 5:143-149.
- Vernet, J.P., and P.Y. Favarger. 1982. Climatic and anthropogenic effects on the sedimentation and geochemistry of Lakes Bourget, Annecy, and Leman. *Hydrobiologia* 92:643-650.
- Vernet, J.P., P.Y. Favarger, and C. Reynaud. 1978. Sedimentation rates in Lake Morat, Joux and Leman computed by means of both ^{137}Cs and pollen-analysis methods. *Intecol, Second Int. Congr. of Ecology, Jerusalem.*
- Villar, H.P. 1981. Distribuicao de ^{137}Cs no solo como funcao de erosao e outros processos. Centro De Energia Nuclear Rep. INIS-MF-7374. Recife, Brazil (In Portuguese.)
- Volchok, H.L., and N. Chieco (ed.) 1986. A compendium of the Environmental Measurements Laboratory's research projects related to the Chernobyl nuclear accident. USDOE Rep. EML-460. EML, New York.
- Von Damm, K.L., L.K. Benninger, and K.K. Turekian. 1979. The ^{210}Pb chronology of a core from Mirror Lake, New Hampshire. *Limnol. Oceanogr.* 24:434-439.
- von Gunten, H.R., M. Sturm, H.N. Erten, E. Rossler, and F. Wegmuller. 1987. Sedimentation rates in the central Lake Constance determined with ^{210}Pb and ^{137}Cs . *Schweiz. Z. Hydrol.* 49:275-283.
- Wahlen, M., and R.C. Thompson. 1980. Pollution records from

- sediments of three lakes in New York State. *Geochim. Cosmochim. Acta* 44:333-339.
- Wahlgren, M.A., and D.M. Nelson. 1975. Plutonium in the Laurentian Great Lakes: Comparison of surface waters. *Verh. Int. Verein. Limnol.* 19:317-322.
- Walling, D.E. 1984. Sediment delivery from drainage basins. p. 71-80. *In* R.J. Loughran (ed.) *Drainage basin erosion and sedimentation*. Univ. of Newcastle, and Soil Conserv. Service of N.S.W., Australia.
- Walling, D.E. 1988. Measuring sediment yield from river basins. p. 39-73. *In* R. Lal (ed.) *Soil erosion research methods*. Soil and Water Conserv. Soc., Ankeny, IA.
- Walling, D.E., and S.B. Bradley. 1988a. The use of caesium-137 measurements to investigate sediment delivery from cultivated areas in Devon, U.K. *IAHS Publ.* 174:325-335.
- Walling, D.E., and S.B. Bradley. 1988b. Transport and redistribution of Chernobyl fallout radionuclides by fluvial processes: Some preliminary evidence. *Environ. Geochem. Health* 10:35-39.
- †Walling, D.E., S.B. Bradley, and C.P. Lambert. 1986a. Conveyance losses of suspended sediment within a flood plain system. p. 119-131. *In* *Drainage basin sediment delivery*. Proc. of the Albuquerque Symp., IAHS Publ. no. 159, Albuquerque, NM. 4-8 August. IAHS Press, Wallingford, U.K.
- †Walling, D.E., S.B. Bradley, and C.J. Wilkinson. 1986b. A caesium-137 budget approach to the investigation of sediment delivery from a small agricultural drainage basin in Devon, U.K. p. 423-435. *In* *Drainage basin sediment delivery*. Proc. of the Albuquerque Symp., IAHS Publ. no. 159, Albuquerque, NM. 4-8 August. IAHS Press, Wallingford, U.K.
- Walling, D.E., and P. Kane. 1984. Suspended sediment properties and their geomorphological significance. p. 311-334. *In* T.P. Burt and D.E. Walling (ed.) *Catchment experiments in fluvial geomorphology*. Geo Books, Norwich, U.K.
- Wan, G., and P.H. Santschi. 1987. Prediction of radionuclide inventory for sediments in Lake Greifen, Switzerland. *Sci. Geogr. Sin.* 4:358-363.
- Wan, G., P.H. Santschi, K. Farrenkoth, M. Sturm, and W. Stumm. 1985. Distribution and dating of ^{137}Cs for recent sediments in Lake Greifen (Switzerland). *Acta Sci. Circumstantia* 5:360-365.
- Wan, G., P.H. Santschi, M. Sturm, K. Farrenkoth, A. Lueck, E. Werth, and C. Schuler. 1986. A comparative study on recent sedimentation rates in Lake Greifen, Switzerland, using varve counting and radionuclide dating. *Geochimica* 9:259-270.
- Wan, G.J., P.H. Santschi, M. Sturm, K. Farrenkoth, A. Lueck, E. Werth, and C. Schuler. 1987. Natural (^{210}Pb , ^7Be) and fallout (^{137}Cs , $^{239,240}\text{Pu}$, ^{90}Sr) radionuclides as geochemical tracers of sedimentation in Greifensee, Switzerland. *Chem. Geol.* 63:181-196.
- Ward, T.E., J. Breeden, K. Komisarik, R. Porter, J. Czuczwa, R. Kaminski, and B.D. McVeety. 1987. ^{137}Cs radioactive dating of Lake Ontario sediment cores. Brookhaven Natl. Lab. Rep. BNL-40550. Brookhaven Natl. Lab., Upton, NY.
- Wasson, R.J., R.L. Clark, P.M. Nanninga, and J. Waters. 1987. ^{210}Pb as a chronometer and tracer, Burinjack Reservoir, Australia. *Earth Surface Processes Landforms* 12:399-414.
- Wasson, R.J., R.L. Clark, I.R. Willett, J. Waters, B.L. Campbell, and D. Outhet. 1984. Erosion history from sedimentation in Burinjack Reservoir, N.S.W. p. 221-229. *In* R.J. Loughran (ed.) *Drainage basin erosion and sedimentation*. Univ. of Newcastle, and Soil Conserv. Service of N.S.W., Australia.
- Wilkin, D.C., and S.J. Hebel. 1982. Erosion, redeposition, and delivery of sediment to Midwestern streams. *Water Resour. Res.* 18:1278-1282.
- Wilken, R.D., I. Moreira, and A. Rebello. 1986. ^{210}Pb and ^{137}Cs fluxes in a sediment core from Guanabara Bay, Brazil. *Sci. Total Environ.* 58:195-198.
- Wilson, G.C., A.E. Litherland, and J.C. Rucklidge. 1984. Dating of sediments using accelerator mass spectrometry. *Chem. Geol.* 44:1-17.
- Wilson, T.R.S. 1974. Caesium-137 as a water movement tracer in the St. George's Channel. *Nature (London)* 248:125-127.
- †Wischmeier, W.H., and D.D. Smith. 1978. Predicting rainfall erosion losses—A guide to conservation planning. *Agric. Handb.* 537 USDA-SEA in cooperation with Purdue Agric. Exp. Stn.
- †Wise, S.M. 1977. The use of fallout radionuclides Pb-210 and Cs-137 in estimating denudation rates and in soil erosion measurement. Occasional Pap. 7, p. 39. Geography Dep., King's College, London.
- †Wise, S.M. 1980. Caesium-137 and lead-210: A review of techniques and some applications in geomorphology. p. 109-127. *In* R.A. Cullingford et al. (ed.) *Timescales in geomorphology*. John Wiley & Sons, New York.
- Woolridge, D.D. 1965. Tracing soil particle movement with Fe-59. *Soil Sci. Soc. Am. Proc.* 29:469-472.
- Wong, G.T.F., and C.S. Moy. 1984. Cesium-137, metals and organic carbon in sediments of the James River estuary, Virginia. *Estuarine Coastal Shelf Sci.* 18:37-49.
- Wong, H.K.T., J.O. Nriagu, and R.D. Coker. 1984. Atmospheric input of heavy metals chronicled in lake sediments of the Algonquin Provincial Park, Ontario, Canada. *Chem. Geol.* 44:187-201.
- Worsley, A.T., and F. Oldfield. 1988. Paleocological studies of three lakes in the highlands of Papua, New Guinea, II. Vegetational history over the last 1600 years. *J. Ecol.* 76:1-18.
- Wrenn, M.E., J.W. Lentsch, M. Eisenbud, G.J. Lauer, and G.P. Howells. 1971. Radiocesium distribution in water, sediment, and biota in the Hudson River estuary from 1964 through 1970. *Proc. Third Natl. Symp. on Radioecology*. USAEC, Oak Ridge, TN.
- Zucker, C.L., C.R. Olsen, I.L. Larsen, and N.H. Cutshall. 1984. Inventories and sorption-desorption trends in radiocesium and radiocobalt in James River estuary sediments. *Environ. Geol. Water Sci.* 6:171-182.